



AN OVERVIEW ON SYNTHETIC METHODS OF N-BUTYL ACETATE

Cong Yufeng^{[a]*}

Keywords: overview; synthetic study; n-butyl acetate; catalysts

Nowadays a few synthetic methods of n-butyl acetate using different catalysts such as inorganic salt like $(\text{Ce}(\text{S}_2\text{O}_8)_2, \text{FeNH}_4(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}, \text{LaSO}_4/\text{SiO}_2, \text{KHSO}_4$ and SnCl_4/C), HZSM-5, oxide ($\text{MoO}_3/\text{SiO}_2$), I_2 , heteropolyacid ($\text{H}_3(\text{PW}_{12}\text{O}_{40}) \cdot n\text{H}_2\text{O}$), quaternary ammonium salt ionic liquid and nanometer ZnO have been reviewed. The yields of n-butyl acetate are improved by the addition of above catalysts. These methods are having the advantages of simple process and low investment costs.

* Corresponding Author

Fax: 86-24-56860869

E-Mail: congyufeng2012@hotmail.com

[a] Liaoning Shihua University, Fushun, Liaoning, P.R. China.

Introduction

n-Butyl acetate is one of the colourless flammable liquids. It is used as a synthetic flavour agent in food products and/or as a solvent in the production of lacquers, plastic or artificial leather.¹ n-Butyl acetate and other organic solvent such as alcohol, aldehyde, ester, etc. are completely miscible in all proportions. Natural n-butyl acetate exists in apple, banana, cherry and grape fruits, etc. It is easily evaporated and hard to dissolve in water. Furthermore, it can dissolve brain grease, gum and rosin. It has anaesthetic and pungent in character. Its specific gravity d_4^{20} , refractive index n_D^{20} and boiling point are 0.8825, 1.3941 and 126.1 °C, respectively.²

In the present paper, different catalysts such as inorganic salt ($\text{Ce}(\text{S}_2\text{O}_8)_2, \text{NH}_4\text{Fe}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}, \text{LaSO}_4/\text{SiO}_2, \text{KHSO}_4$ and SnCl_4/C), HZSM-5, oxide ($\text{MoO}_3/\text{SiO}_2$), I_2 , heteropolyacid ($\text{H}_3(\text{PW}_{12}\text{O}_{40}) \cdot n\text{H}_2\text{O}$), quaternary ammonium salt ionic liquid and nanometer ZnO have been discussed.

DISCUSSION

$\text{Ce}(\text{S}_2\text{O}_8)_2$ -SBA-15 as a catalyst to produce n-butyl acetate

Yin Yanlei³ introduced the preparation of n-butyl acetate and the effect of the reaction conditions on its yield. $\text{Ce}(\text{S}_2\text{O}_8)_2$ and SBA-15 as catalysts were ground and roasted to produce n-butyl acetate. The optimum conditions were the reaction time (2 hours) and molar ratio of acetic acid to n-butanol (1.0:1.2), amount of catalyst (0.0375g) respectively. The maximum yield of n-butyl acetate was 95.13%. The experimental results showed that $\text{Ce}(\text{S}_2\text{O}_8)_2$ and SBA-15 had good catalytic performance and were reused several times.

$\text{NH}_4\text{Fe}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ as a catalyst to generate n-butyl acetate

Kong Xiangwen⁴ explained why $\text{NH}_4\text{Fe}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ as a catalyst took the place of concentrated sulfuric acid to generate

n-butyl acetate. The effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst had been discussed. The experimental results represented that the best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 0.75 hours, 1.0:1.2 and 1.1 g, respectively. The maximum yield of n-butyl acetate was 98.5%.

$\text{LaSO}_4/\text{SiO}_2$ as a catalyst to produce n-butyl acetate

Song Jianguo⁵ described the synthetic method of n-butyl acetate by using $\text{LaSO}_4/\text{SiO}_2$ as catalysts. The effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol, amount of catalyst and number of reusable catalyst had been discussed. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 2 hours, 1.0:1.3 and 0.06 g, respectively. The maximum yield of n-butyl acetate was 95.2%. After recovery and reuse of $\text{LaSO}_4/\text{SiO}_2$ as a catalyst it was noticed that its catalytic performance was very poor. For example, when $\text{LaSO}_4/\text{SiO}_2$ were reused 5 times, the yield of n-butyl acetate only reached 56.3%.

KHSO_4 as a catalyst to generate n-butyl acetate

Cai Xin'an⁶ used KHSO_4 as a catalyst to produce n-butyl acetate. The effect of the reaction conditions such as amount of n-butanol, molar ratio of acetic acid to n-butanol and amount of catalyst had been discussed. The best conditions were that amount of n-butanol, molar ratio of acetic acid to n-butanol and amount of catalyst were 0.25 mol, 1.0:1.3 and 1.15 g, respectively. The maximum yield of n-butyl acetate was 86.83%.

SnCl_4/C as a catalyst to produce n-butyl acetate

Li Jiagui⁷ described the synthetic method of n-butyl acetate by using SnCl_4/C as catalysts. The effect of the reaction conditions such as amount of acetic acid, molar ratio of acetic acid to n-butanol, amount of catalyst and number of reusable catalyst had been discussed. The best conditions were that amount of acetic acid, molar ratio of acetic acid to n-butanol and amount of catalyst were 0.1 mol, 1.0:2.2 and 0.15 g, respectively. The maximum yield of n-butyl acetate was 94.8%. When SnCl_4/C as catalysts were recovered and reused, their catalytic

performance was found to be very good. For example, when SnCl₄/C were reused 5 times, the yield of n-butyl acetate still reached 90.1%.

HZSM-5 as a catalyst to generate n-butyl acetate

Li Minghui⁸ introduced the preparation of HZSM-5 and the effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst on the yield of n-butyl acetate. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 1 hour, 1.0:1.1 and 0.5 g, respectively. The maximum yield of n-butyl acetate was 94.0%.

MoO₃/SiO₂ as a catalyst to produce n-butyl acetate

Li Shuchang⁹ described the preparation of MoO₃/SiO₂ and the effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol, amount of catalyst and number of reusable catalyst on the yield of n-butyl acetate. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 3 hours, 1.0:4.0 and 1.0 g, respectively. The maximum yield of n-butyl acetate was 95.6%. When MoO₃/SiO₂ as catalysts were recovered and reused, their catalytic performance was perfect. For example, when MoO₃/SiO₂ were reused 5 times, the yield of n-butyl acetate still arrived at 95.2%.

I₂ as a catalyst to generate n-butyl acetate

Jiang Hongbo¹⁰ used I₂ as a catalyst to generate n-butyl acetate. The effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst had been discussed. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 4.0 hours, 1.5:1.0 and 0.6 g, respectively. The maximum yield of n-butyl acetate was 67.52%.

H₂(PW₁₂O₄₀)·nH₂O as a catalyst to produce n-butyl acetate

Li Guixian¹¹ introduced the preparation of H₃(PW₁₂O₄₀)·nH₂O with attapulgite and the effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol, amount of catalyst and number of usable catalyst on the yield of n-butyl acetate. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 2.5 hours, 1.0:2.5 and 0.1580 g, respectively. The maximum yield of n-butyl acetate was 94.0%. When H₃(PW₁₂O₄₀)·nH₂O as a catalyst was recovered and reused, its catalytic performance was poor. For example, after used 5 times as catalyst, H₃(PW₁₂O₄₀)·nH₂O, the yield of n-butyl acetate reached only to 77.98%.

Quaternary ammonium salt ionic liquid as a catalyst to generate n-butyl acetate

Li Wenjun¹² described the preparation of four types of quaternary ammonium salt ionic liquids such as [Et₂NH₂][HSO₄], [Et₃NH][HSO₄], [n-PrO₃NH][HSO₄] and [n-

But₃NH][HSO₄]. He used [Et₃NH][HSO₄] as a catalyst to generate n-butyl acetate. The effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol and molar ratio of [Et₃NH][HSO₄] to n-butanol had been discussed. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and molar ratio of [Et₃NH][HSO₄] to n-butanol were 8.0 hours, 2.0:1.0 and 1.0:4.0, respectively. The maximum yield of n-butyl acetate was 81.94%.

Nanometer ZnO as a catalyst to produce n-butyl acetate

Lai Wenzhong¹³ used nanometer ZnO as a catalyst and acetic acid to n-butanol as feedstocks to produce n-butyl acetate. The effect of the reaction conditions such as the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst on the yield of n-butyl acetate had been discussed. The best conditions were that the reaction time, molar ratio of acetic acid to n-butanol and amount of catalyst were 3.0 hours, 1.0:2.0 and 0.80 g, respectively. The maximum yield of n-butyl acetate was 86.72%.

CONCLUSION

Based on the above discussion and review, NH₄Fe(SO₄)₂·12H₂O that is one of the best catalysts is used to produce n-butyl acetate because it is very cheap, stable and insoluble in organic acids and organic alcohol. After the reaction is done, NH₄Fe(SO₄)₂·12H₂O becomes insoluble material and is easily separated from the reaction system. It has high performance catalytic and selectivity and non-corrosive. However, I₂ is one of the worst catalysts and its maximum yield of n-butyl acetate only reaches 67.52%.

REFERENCES

- ¹An, J. J. and Wang, B. Y. *Beijing Light Industry Press*, **1989**.
- ²You, H. J. *Sichuang Chem. Ind.*, **2005**, 8(1), 15.
- ³Yin, Y. L., Tian, Z. M., Deng, Q. G. and Ding, K. P. *J. Qiqihar Univ.*, **2008**, 24(1), 4.
- ⁴Kong, X. W., Yu, L., Zhang, J. and Shu, Y. *Res. Exploration Lab.*, **2011**, 30(2), 21.
- ⁵Song, J. G. *J. Qiqihar Univ.*, **2008**, 24(3), 47.
- ⁶Cai, X. A., Chen, L. X., Zhang, H. F., Wang, L. and Wang, K. *K. J. Jingdezhen College*, **2012**, 27(3), 37.
- ⁷Li, J. G., Wei, Q. M. and Zhu, W. R. *J. Yulin Normal Univ.*, **2008**, 29(5), 47.
- ⁸Li, M. H., Zhu, M. and Liu, N. *J. Dalian Polytechnic Univ.*, **2009**, 28(5), 347.
- ⁹Li, S. C., Sun, D. and Du, C. H. *Chem. Eng.*, **2010**, 10, 61.
- ¹⁰Jiang, H. B. *Chem. Eng.*, **2011**, 6, 14.
- ¹¹Li, G. X., Wang, X. N., Shi, A. J., Wang, K. Z. and Hu, D. C. *J. Mol. Catal.*, (*Fenzi Cuihua*) **2008**, 22(2), 137.
- ¹²Li, W. J., Zhong, L. N., Chen, Y. Q. and Lin, J. Q. *J. Chem. Ind. Eng.*, **2011**, 62(S2), 74.
- ¹³Lai, W. Z., Lin, Y. and Xiao, W. C. *Food Sci. Technol.*, **2010**, 35(2), 210.

Received: 22.10.2012.
Accepted: 24.10.2012.