



SILICA SUPPORTED PERCHLORIC ACID: AN EFFICIENT AND RECYCLABLE CATALYST FOR SYNTHESIS OF BENZIMIDAZOLO[2,3-*b*]QUINAZOLINONES

Vinod V. Throat,^[a] Maya V. Katariya,^[a] Sunil U. Tekale,^[a] Rupali L. Magar,^[a] Samson Mashele,^[b] Pravin S. Kendrekar^{[b]*} and Rajendra P. Pawar^{[b]*}

Keywords: 2-Aminobenzimidazole; benzimidazolo[2,3-*b*]quinazolinones; heterogeneous catalyst; perchloric acid; silica.

Synthesis of benzimidazolo[2,3-*b*]quinazolinone derivatives has been reported in excellent yields by using silica-supported perchloric acid (HClO₄-SiO₂) as a mild and reusable heterogeneous catalyst. The procedure is simple, environmentally benign and has the advantage of high atom economy. Furthermore, the catalyst can be recovered and reused several times efficiently without substantial loss of catalytic activity.

*Corresponding author

Tel: +91 0240 2334577; Fax: +91 0240- 2334430

E-Mail: rppawar@yahoo.com, kkpravin@gmail.com

[a] Department of Chemistry, Deogiri College, Aurangabad, 431005, MS, India.

[b] Department of Health Sciences, Central University of Technology, Free State, South Africa.




Figure 1. Biologically important quinazolines


INTRODUCTION

Heterogeneous catalysis is an interesting area of research from an industrial point of view. It has the advantages of thermal stability, high selectivity, better activity, ease of separation, recyclability and long life.¹⁻⁴ Solid acid catalysts play an important role in organic transformations due to many advantages such as simplicity in handling, decreased reactor corrosion problems and more environmentally safe disposal of the catalyst.⁵⁻⁷ Quinazolinones and their derivatives have been reported to possess interesting pharmacological activities, such as antibacterial,⁸⁻⁹ antihypertension,¹⁰ antihistaminic,¹¹ analgesic, anti-inflammatory,¹² anticancer,¹³ and anti-HIV.¹⁴ Moreover, a variety of quinazolinones derivatives with different biological activities were synthesized by medicinal chemistry researchers. These derivatives also have a long history of applications in agrochemicals and the pharmaceutical industry as herbicides and active pharmaceuticals. Awareness about environmental hazards in chemical industries becomes a significant concern due to the generation of waste products that leads to the development of environment-friendly synthetic processes. Heterocyclic compounds constitute comprehensive examples in pharmaceutical and chemical industries. Because of their potent physiological properties, they resulted in numerous applications.¹⁵

Several methods have been reported for the synthesis of substituted benzimidazolo-quinazolinones. The most common method is the reaction of substituted aldehydes with 2-aminobenzimidazole and dimedone using various basic and acidic catalysts under reflux conditions,¹⁵⁻¹⁶ ionic liquids¹⁷ and heteropolyacids.¹⁸

All these methods are associated with several limitations such as the use of metal catalysts, harsh reaction conditions, tedious experimental procedure, low yields, prolonged reaction time and use of costly and moisture-sensitive catalyst. Hence, there is a need to develop a rapid, efficient and environmentally benign synthetic procedure for the synthesis of benzimidazole[2,3-*b*]quinazolinone derivatives.

In continuation of previous studies on silica-supported perchloric acid¹⁹⁻²¹ herein, we developed an efficient and environment-friendly method for the synthesis of quinazolinone derivatives by the condensation of substituted aldehydes, 2-aminobenzimidazole, and dimedone using silica-supported perchloric acid as a heterogeneous catalyst (Scheme 1).



Scheme 1. Synthesis of benzimidazolo[2,3-*b*]quinazolinones

EXPERIMENTAL □

Preparation of HClO₄-SiO₂ catalyst:

Aqueous perchloric acid (70 %, 1.8g, 12.5 mmol) was added to a suspension of SiO₂ (230-400 mesh, 23.7 g) in

ether (70 ml). The mixture was concentrated and the residue was heated at 100 °C for 72 h under vacuum to give HClO₄-SiO₂ (0.5 mmol g⁻¹) as free-flowing powder.²²

General procedure: synthesis of 3,3-dimethyl-12-phenyl-3,4,5,12-tetrahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1(2*H*)-one

Silica supported perchloric acid (10 wt.%) was added to a mixture of 2-aminobenzimidazole (1 mmol), aldehyde (1 mmol) and dimedone (1) mmol in 1:1 ethanol: water (5 mL). The reaction mixture was stirred at 80 °C for 20-40 minutes. After completion of the reaction, as monitored by TLC, the reaction mass was filtered. The filtrate was heated to remove the solvent. Separated solid was washed with water and dried under reduced pressure. Furthermore, the separated catalyst was dried and reused.

Similarly, the other derivatives were also synthesized using the same method (Table 1). Spectral data of the synthesized compounds is mentioned below:

3,3-Dimethyl-12-phenyl-1,2,3,4,5,12-hexahydrobenzo[4,5]-imidazo[2,1-*b*]quinazolin-1-one (4a)

M.p. 270-280 °C; IR (KBr): 3350, 2920, 1640, 1620, 1610, 1565 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δH = 1.06 (s, 3H, CH₃), 1.09 (s, 3H, CH₃), 2.04 (s, 2H, CH₂), 2.30 (s, 2H, CH₂), 6.41 (s, 1H, CH), 6.93-7.38 (m, 9H, Ar-H), 11.12 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(2,4-dichlorophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4b)

M.p. 315-330 °C; IR (KBr): 3085, 2930, 1615, 1520 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.0 (s, 3H, CH₃), 1.08 (s, 3H, CH₃), 2.19 (s, 2H, CH₂), 2.35 (s, 2H, CH₂), 6.90 (s, 1H, CH), 7.25-8.10 (m, 4H, Ar-H), 7.25-8.10 (m, 2H, Ar-H), 8.15 (s, 1H, Ar-H), 9.11.31 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-bromophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4c)

M.p. 295-300 °C; IR (KBr): 3420, 2920, 1640, 1610, 1530 cm⁻¹; ¹H NMR (DMSO-*d*₆, 300 MHz): δ H = 0.90 (s, 3H, CH₃), 1.05 (s, 3H, CH₃), 2.00 (s, 2H, CH₂), 2.20 (s, 2H, CH₂), 6.42 (s, 1H, CH), 6.98-7.88 (m, 8H, Ar-H), 11.00 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-nitrophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4d)

M.p. 290-300 °C; IR (KBr): 2869, 1681, 1612, 1518 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 0.85 (s, 3H, CH₃), 1.6 (s, 3H, CH₃), 2.20 (s, 2H, CH₂), 2.28 (s, 2H, CH₂), 6.60 (s, 1H, CH), 7.04-8.10 (m, 8H, Ar-H), 11.90 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-fluorophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4e)

M.p. 285-295 °C; IR (KBr): 3020, 2915, 1645, 1580, 1330 cm⁻¹; ¹H NMR (DMSO-*d*₆, 300 MHz): δ H = 1.04 (s, 3H, CH₃), 1.07 (s, 3H, CH₃), 2.20 (s, 2H, CH₂), 2.04 (s, 2H, CH₂), 6.49 (s, 1H, CH), 6.90-7.90 (m, 8H, Ar-H), 10.90 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-chlorophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4f)

M.p. 285-295 °C; IR (KBr): ν_{max} = 3465, 2945, 1670, 1619, 1560, 1566 cm⁻¹; ¹H NMR (DMSO-*d*₆, 300 MHz): δ H = 1.06 (s, 3H, CH₃), 1.11 (s, 3H, CH₃), 2.03 (s, 2H, CH₂), 2.40 (s, 2H, CH₂), 6.56 (s, 1H, CH), 6.94-7.53 (m, 8H, Ar-H), 11.27 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(2,4,6-methoxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4g)

M.p. 292-302 °C; IR (KBr): 3210, 2969, 1690, 1590, 1312, 1258 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.09 (s, 3H, CH₃), 1.16 (s, 3H, CH₃), 2.11 (s, 2H, CH₂), 2.49 (s, 2H, CH₂), 3.69 (s, 3H, OCH₃), 3.75 (s, 6H, OCH₃), 6.30 (s, 1H, CH), 6.55 (s, 2H, Ar-H), 6.73-7.42 (m, 4H, Ar-H), 11.01 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-methoxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4h)

M.p. 280-290 °C; IR (KBr): 3243, 2961, 1680, 1641, 1612, 1589, 1566, 1258 cm⁻¹; ¹H NMR (DMSO-*d*₆, 300 MHz): δ H = 1.03 (s, 3H, CH₃), 1.06 (s, 3H, CH₃), 2.06-2.15 (m, 2H, CH₂), 2.25-2.50 (m, 2H, CH₂), 3.66 (s, 3H, OCH₃), 6.35 (s, 1H, H-12), 6.73-7.42 (m, 8H, Ar-H), 11.01 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(2,4-methoxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4i)

M.p. 250-260 °C; IR (KBr): 3085, 2925, 1600, 1575 1262 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.06 (s, 3H, CH₃), 1.11 (s, 3H, CH₃), 1.90 (s, 2H, CH₂), 2.45 (s, 2H, CH₂), 3.72 (s, 3H, OCH₃), 3.83 (s, 3H, OCH₃), 6.20 (s, 1H, CH), 6.54-7.44 (m, 3H, Ar-H), 6.58 (t, 1H, Ar-H), 7.44 (d, 1H, Ar-H), 7.00-7.50 (m, 4H, Ar-H), 11.21 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-hydroxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4j)

M.p. 270-275 °C; IR (KBr): 3469, 2962, 1574, 1264 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.06 (s, 3H, CH₃), 1.06 (s, 3H, CH₃), 2.04 (s, 2H, CH₂), 2.43 (s, 2H, CH₂), 6.19 (s, 1H, CH), 6.60-7.35 (m, 8H, Ar-H), 8-9.32 (s, 1H, OH), 11.01 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(3-hydroxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4k)

M.p. 287-292 °C; IR (KBr): 3090, 2950, 1572, 1249 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.07 (s, 3H, CH₃), 1.09 (s, 3H, CH₃), 1.95 (s, 2H, CH₂), 2.50 (s, 2H, CH₂), 6.23 (s, 1H, CH), 6.69-7.00 (m, 3H, Ar-H), 7.10 (t, J = 7.90, Hz, 1H, Ar-H), 7.52 (d, J = 5.69, Hz, 1H, Ar-H), 7.30-7.70 (m, 4H, Ar-H), 9.20 (s, 1H, OH), 11.30 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(4-methylphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4l)

M.p. 260-270 °C; IR (KBr): 3085, 2930, 1570, 1253 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.10 (s, 3H, CH₃), 1.12 (s, 3H, CH₃), 2.15 (s, 2H, CH₂), 2.40 (s, 2H, CH₂), 2.70 (s, 3H, CH₃), 6.35 (s, 1H, H-12), 6.60-7.35 (m, 8H, Ar-H), 11.01 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(2-nitrophenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4m)

M.p. 275-280 °C; IR (KBr): 3400, 2995, 1589, 1320 1258, 749 cm⁻¹; ¹H NMR (DMSO-*d*₆, 300 MHz): δ H = 1.07 (s, 3H, CH₃), 1.11 (s, 3H, CH₃), 2.19 (s, 2H, CH₂), 2.70 (s, 2H, CH₂), 6.45 (s, 1H, CH), 7.05 (d, 1H, Ar-H), 7.20-7.35 (m, 1H, Ar-H), 7.15-7.44 (m, 1H, Ar-H), 7.15-7.44 (m, 1H, Ar-H), 7.44 (d, J = 8.15 Hz, 1H, Ar-H), 7.50-7.90 (m, 4H, Ar-H), 11.29 (s, 1H, NH) ppm.

3,3-Dimethyl-12-(2-methoxyphenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (4n)















M.p. 245-255 °C; IR (KBr): 3089, 2895, 1590, 1248, 740 cm⁻¹; ¹H-NMR (DMSO-*d*₆, 300 MHz): δ H = 1.09 (s, 3H, CH₃), 1.14 (s, 3H, CH₃), 1.99 (s, 2H, CH₂), 2.45 (s, 2H, CH₂), 3.72 (s, 3H, OCH₃), 6.29 (s, 1H, CH), 6.90-7.44 (m, 4H, Ar-H), 7.15-7.64 (m, 4H, Ar-H), 11.27 (s, 1H, NH) ppm.

RESULTS AND DISCUSSION

We developed a new method for the synthesis of benzimidazolo[2,3-*b*]quinazolinone derivatives in good yields by using HClO₄-SiO₂ as a mild and reusable heterogeneous catalyst in water:ethanol (1:1) solvent. The procedure is environment-friendly, operationally simple and thus has the advantage of high atom economy. Furthermore, the catalyst can be recovered and reused several times efficiently without substantial loss of activity. We also studied the reaction in aqueous medium only, but the reaction proceeded very slowly and the product formation was also very poor. When we used ethanol:water (1:1) solvent system, the reaction proceeded faster with a high yield of the corresponding product.

The effect of temperature on yield of the product **4a** was studied by carrying the reactions at different temperatures (30, 55, 80 and 100 °C) as shown in Table 2. By raising the reaction temperature from room temperature to 100 °C gradually, the yield of reactions was found to be increased.

Table 1. Synthesis of 3,3-dimethyl-12-(un)substituted phenyl-3,4,5,12-tetrahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1(2*H*)-ones using silica supported perchloric acid.

Entry	Aldehydes (2)	Products (4)	Time (min)	Yield (%)
1		4a	30	85
2		4b	30	90
3		4c	40	85
4		4d	20	80
5		4e	25	75
6		4f	25	90
7		4g	30	80
8		4h	20	90
9		4i	25	90
10		4j	35	75
11		4k	30	90
12		4l	20	90
13		4m	25	90
14		4n	20	90

Reaction conditions: Aldehyde (1.0 mmol), dimedone (1.0 mmol), 2-aminobenzimidazole (1.0 mmol), HClO₄-SiO₂ (10 wt%) 20-40 min reflux.

At 80 °C temperature, the reaction completed in 25 minutes affording 90 % of product yield. Similarly, increasing the reaction temperature to 100 °C does not affect the yield of the product significantly. Thus, we confirmed that 80 °C was the optimum temperature for the transformation. Under these optimized conditions, various aldehydes were reacted with dimedone and aminobenzimidazoles, whose results are summarized in Table 1.

Table 2. Effect of temperature on the preparation of 3,3-dimethyl-12-phenyl-1,2,3,4,5,12-hexahydrobenzo[4,5]-imidazo[2,1-*b*]quinazolin-1-one (**4a**)

Entry	Temp., °C	Time, min	Yield, % [@]
1	30	240	40
2	50	60	70
3	80	25	90
4	100	25	90

[@]Reactions performed in case of 4-chlorobenzaldehyde

Reusability of the catalyst

Solid silica-based perchloric acid works under heterogeneous conditions. It is an inexpensive and non-hazardous solid acid catalyst which can be easily handled and separated from the reaction mixture by simple filtration. The recovered catalyst was reused thrice for consecutive runs with a minimum variation of yields of the products. After completion of the reaction, the catalyst was filtered, thoroughly washed with ethanol and dried at 100 °C for 2 hr and reused for subsequent runs (Table 3). This reusability demonstrates the high stability and turnover of solid silica-based perchloric acid under operating conditions.

Table 3. Reusability of catalyst for the synthesis of 3,3-dimethyl-12-phenyl-1,2,3,4,5,12-hexahydrobenzo[4,5]-imidazo[2,1-*b*]quinazolin-1-one (**4a**)

Entry	Number of recycling □	Time, min	Yield, % [@]
1	Fresh	40	90
2	First	45	88
3	Second time	60	88
4	Third time	70	85

[@]Yields in case of 4-chlorobenzaldehyde

CONCLUSION

A convenient method has been developed by the reaction of 2-aminobenzimidazole, dimedone and aldehyde catalyzed by the silica-supported perchloric acid catalyst. Use of inexpensive and reusable catalyst, enhanced reaction rates, readily available starting materials, high yield and easy purification of the products are the key features of this method.

ACKNOWLEDGMENTS

The authors are thankful to the Principal, Deogiri College, Aurangabad (MS), India for encouragement and providing laboratory facilities.

REFERENCES

- ¹Choudhary, D., Paul, S., Gupta, R., Clark, J. H., Catalytic properties of several palladium complexes covalently anchored on to silica for the aerobic oxidation of alcohols, *Green Chem.*, **2006**, *8*, 479-482. DOI:10.1039/b601363g
- ²Li, Z., Ma, X., Liu, J., Feng, X., Tian, G., Zhu, A., Silica-supported aluminum chloride: A recyclable and reusable catalyst for one-pot three-component Mannich-type reactions, *J. Mol. Catal. A: Chem.*, **2007**, *272*, 132-135. <https://doi.org/10.1016/j.molcata.2007.03.029>
- ³Hongjin, L., Song, J., Geletii, Y., Vickers, J., Sumliner, J., An exceptionally fast homogeneous carbon-free cobalt-based water oxidation catalyst, *J. Am. Chem. Soc.*, **2014**, *136*, 9268. <https://pubs.acs.org/doi/10.1021/ja5045488>
- ⁴Melero, J. A., Grieken, R.V., Morales, G., Advances in the synthesis and catalytic applications of organosulfonic-functionalized mesostructured materials, *Chem Rev.*, **2006**, *106*, 3790. <https://doi.org/10.1021/cr050994h>
- ⁵Niknam, K., Saberi, D., Silica-bonded *N*-propyl sulfamic acid as an efficient catalyst for the formylation and acetylation of alcohols and amines under heterogeneous conditions, *Tetrahedron Lett.*, **2009**, *50*, 5210. <https://doi.org/10.1016/j.tetlet.2009.06.140>
- ⁶Niknam, K., Saberi, D., Nouri, S.M., Silica-bonded *S*-sulfonic acid as a recyclable catalyst for chemoselective synthesis of 1,1-diacetates, *Tetrahedron Lett.*, **2009**, *50*, 4058. <https://doi.org/10.1016/j.tetlet.2009.04.096>
- ⁷Niknam, K., Zolfigol, M. A., Sadabadi, T., Ca(HSO₄)₂ mediated conversion of alcohols into *N*-substituted amides under heterogeneous conditions: A modified Ritter reaction, *J. Iran. Chem. Soc.*, **2007**, *4*, 199. <https://doi.org/10.1007/BF03245967>
- ⁸Ozden, S., Atabey, D., Yıldız, S., Göker, H., Synthesis and potent antimicrobial activity of some novel methyl or ethyl 1*H*-benzimidazole-5-carboxylates derivatives carrying amide or amidine groups, *Bioorg. Med. Chem.*, **2005**, *13*, 1587. <https://doi.org/10.1016/j.bmc.2004.12.025>
- ⁹Ansari, K. F., Lal, C., Synthesis, physicochemical properties and antimicrobial activity of some new benzimidazole derivatives, *Eur. J. Med. Chem.*, **2009**, *44*, 4028. <https://doi.org/10.1016/j.ejmech.2009.04.037>
- ¹⁰Alagersamy, V., Pathak, Synthesis and antihypertensive activity of novel 3-benzyl-2-substituted-3*H*-[1,2,4]triazolo[5,1-*b*]quinazolin-9-ones, *Biorganic Med. Chemistry.*, **2007**, *15*, 3457. <https://doi.org/10.1016/j.bmc.2007.03.007>
- ¹¹Alagersamy, V., Murugan, M., Synthesis and pharmacological investigation of novel 4-benzyl-1-substituted-4*H*-[1,2,4]triazolo[4,3-*a*]quinazolin-5-ones as new class of H₁-antihistaminic agents, *Biorganic Med. Chemistry*, **2007**, *15*, 4009. <https://doi.org/10.1016/j.bmc.2007.04.001>
- ¹²Alagersamy, V., Muruganathan, G., Venkateshperumal, R., Synthesis, Analgesic, Anti-inflammatory and antibacterial activities of some novel 2-methyl-3-substituted quinazolin-4-(3*H*)-ones. *Biol. Pharm. Bull.*, **2003**, *26*, 1711. <https://doi.org/10.1248/bpb.26.1711>
- ¹³Valeriy, A., Bacherikov, T. C., Hua-Jin, D., Potent antitumor *N*-mustard derivatives of 9-anilinoacridine, synthesis and antitumor evaluation, *Biorganic Med. Chemistry*, **2004**, *14*, 4719. <https://doi.org/10.1016/j.bmcl.2004.06.080>

- ¹⁴Kumar, D., Mariappan, G., Husain, A., Monga, J., Kumar, S., Design, synthesis and cytotoxic evaluation of novel imidazolone fused quinazolinone derivatives, *Arab. J. Chem.*, **2017**, *10*, 344. <https://doi.org/10.1016/j.arabjc.2014.07.001>
- ¹⁵Lipson, V. V., Desenko, S. M., Shirobokova, M. G Shishkin, O. V., Orlov, V. D., Cyclocondensation of 2-aminobenzimidazole with dimedone and its arylidene derivatives, *Chem. Het. Com.*, **2003**, *39*, 1041. <https://doi.org/10.1023/B:COHC.000003522.71893.77>
- ¹⁶Kumari, K., Raghuvanshi, D. S., Singh, K. N. An Expeditious Synthesis of Tetrahydro-1,2,4-triazolo[5,1-*b*]quinazolin-8(4*H*)-ones and Dihydro-1,2,4-triazolo[1,5-*a*]pyrimidines, *Org. Prep. Proced. Int.*, **2012**, *44*, 460. <https://doi.org/10.1080/00304948.2012.715062>
- ¹⁷Heravi, M. M., Ranjbar, L., Derikvand, F., Alimadadi, B., Oskooie, H. A., Bamoharram, F. F., A three-component one-pot procedure for the synthesis of [1,2,4]triazolo/benzimidazolo-quinazolinone derivatives in the presence of H₆P₂W₁₈O₆₂ · 18H₂O as a green and reusable catalyst, *Mol. Divers.*, **2008**, *12*, 181-185. DOI [10.1007/s11030-008-9086-8](https://doi.org/10.1007/s11030-008-9086-8)
- ¹⁸Puligoundlaa, R. G., Karnakanti, S., A simple, convenient one-pot synthesis of [1,2,4]triazolo/benzimidazolo quinazolinone derivatives by using molecular iodine, *Tetrahedron Lett.*, **2013**, *54*, 2480. <https://doi.org/10.1016/j.tetlet.2013.02.099>
- ¹⁹Gholam, H. M., Mohammad, A. B., Silica supported perchloric acid (HClO₄-SiO₂): A mild, reusable and highly efficient heterogeneous catalyst for the synthesis of amidoalkyl naphthols *Chin. Chem. Lett.*, **2008**, *19*, 1171. <https://doi.org/10.1016/j.ccllet.2008.06.048>
- ²⁰Hamid, R. S., Majid, G., Silica supported perchloric acid (HClO₄-SiO₂): an efficient and recyclable heterogeneous catalyst for the one-pot synthesis of amidoalkyl naphthols *Tetrahedron*, **2008**, *64*, 1263. <https://doi.org/10.1016/j.tet.2007.11.070>
- ²¹Vinod, K., Vasant, S. J., Neet, J., An efficient method for the synthesis of acylals from aldehydes using silica-supported perchloric acid (HClO₄-SiO₂), *Tetrahedron Lett.*, **2006**, *47*, 5573. <https://doi.org/10.1016/j.tetlet.2006.05.125>
- ²²Babasaheb, P. B., Shrikant, S. G., Dnyanoba, B. M., Silica supported perchloric acid (HClO₄-SiO₂): A green, reusable, and highly efficient heterogeneous catalyst for the synthesis of thioethers under solvent-free conditions at room temperature, *Green Chem. Lett. Rev.*, **2010**, *49*, 54. <https://doi.org/10.1080/17518250903447118>

Received: 04.03.2019.

Accepted: 13.09.2019.