



GASTROPODS AS MARKERS IN BABURA RIVER AT NORTH SUMATRA

Masdiana Sinambela^{1*}, Alex T. Barus², B. Manurung¹, Hesti Wahyuningsih²

¹Department Biology, Faculty Of Mathematics And Natural Sciences, Universitas Negeri Medan,
Medan, Indonesia

²Department Biology, Faculty Of Mathematics And Natural Sciences, Universitas Sumatera Utara,
Medan, Indonesia

*Corresponding Author :

Email: masdiana@unimed.ac.id

doi: 10.48047/ecb/2023.12.1.328

Abstract

Gastropod from the Babura river were taken by hand and an eckman grab. There were 6 research station, (I) in upstream areas, (II) after plantation areas, (III) after pig farm, (IV) after sand mining, (V) after settlements, and (VI) in downstream areas, sampling from March 2017 to January 2018. Gastropod samples were sent for identification to LIPI Cibinong so that the results obtained were more accurate. Diversity, similarity and dominance index of the gastropod spesies were determined. Data were sratically verified by PCA using SAS 9.1.3 package. Gastropod community in the river consisted of : 6 genera, namely, (1) Pomacea, (2) Filopaludina, (3) Lymnaea, (4) Terebia, (5) Melanoides, (6) Thiara and 1 sub-genera, namely Clea (Anentome). Diversity at station 4 is low and at stations 2, 5, and 6 is moderate. Similirity at stations 3 to 6 is high. The dominance at stations 3, 5, and 6 is low and at station 4 is medium. Parameters that greatly affect the presence of each species of gastropods in the Babura River are: *Pomacea canaliculata* by DO, *Thiara scraba* by current flow, *Tarebia granifera* by BOD, *Melanoides tuberculata* by pH, *Lymnaea rubiginosa* by current flow, *Filopaludina javanica* by temperature, and *Clea (Anentome) helena* by current flow. The gastropods markers in the Babura river are *Lymnaea rubiginosa* and *Clea (Anentome) helena*.

Keywords: Markers, dominance, gastropod, correlation and parameters.

Introduction

Gastropods play an important role in an ecosystem because of their ability as a filter feeder that filters dissolved substances in the water (Kushadiwijayanto *et al*, 2022) and can be used as a bioindicator of water quality because it is sensitive to water pollution (Afwanudin *et al*, 2019 & Harahap A., 2021.). *Melanoide tuberculatus* has a high tolerance for environmental changes and is

one of the invasive gastropods (Abdelhady *et al*, 2018). According to Sinambela *et al* (2019), 6 species of gastropods were found in the Babura river.


Babura river water comes from seeps, the upstream is located in the village of Keci-keci Bingkawang Sibolangit forest. The Babura River crosses the area of plantation, pig farms, sand mining, settlements, and downstream in Petisah,

Medan city. The river has a long flow as the source of life by humans, like bathing, washing, and so on as a livelihood (Shinta *et al.*, 2020 & Harahap, A., et all 2022). Currently, there have been many studies conducted on river water with the aim of monitoring and controlling water quality (Satar, 2017). This study aims to determine gastropod as markers in Babura river at North Sumatra.

Gastropod

Gastropods communities are the most diverse class of mollusc and considered as foremost constituent in the freshwater ecosystems (Hamli *et al.*, 2020 & Harahap, A., et all 2022.). The occupied by gastropods have been continuously disrupted by various factors (Jung & Seuk, 2020). Gastropods can be life in the characteristics of waters are sand, muddy and muddy sand (Kushadiwijayanto *et al.*, 2022 & K.Khairul, A Harahap, 2019), gastropods are also found in freshwater (Hamli *et al.*, 2020). Gastropods include macrozoobenthos which can be observed with the naked eye (Moghdani *et al.*, 2013). Ancient lakes in Sulawesi show cases of diversification of freshwater gastropod groups (Albrecht *et al.*, 2020). Gastropods have developed a unique metabolite that acts as a potential competitive slug deterrent, (Raw *et et al.*, 2015; Kushadiwijayanto *et al.*, 2022).

Figure 1. Sampling Research Station on the Babura River



The Babura Watershed (DAS) administratively covers part of Deli Serdang Regency and the Medan city, covering an area of \pm 4921.88 Ha, from upstream in Sibolangit District, Deli Serdang to downstream in Petisah Medan. Its up stream comes from springs and see pages from cliffs so that the water discharge is relatively small. In the middle part of Namorambe district, several sand excavations were found. The lower reaches of the Babura river is where it meets the Deli river (Narayan *et al.*, 2023), (Narayan *et al.*, 2022).

The research stations were detemined based on the environmental baseline, namely station I in the upstream area with coordinate point 3021'22.6"N 98035'34.3"E, station II after people's gardens with coordinate point 3024'13.0"LU 98037'17.9"E, station III after pig farming with coordinate point 3029'48.5"N 98039'11.8"E, station IV after sand mining with coordinate point 3024'52.6"N 98038'52.6"E, station V after settlement with coordinates point 3030'34.9"N 98039'31,5"BT, station VI in the city of Medan downstream area with coordinate point 3034'30.0"N 98040'08.0"BT (Figure 1), sampling from March 2017 to January 2018. Identification of gastropod to the Zoology laboratory of LIPI Cibinong Bogor (Sinambela *et al.*, 2019) and analysis of water samples to BTKL in Medan city (Babu *et al.*, 2020)

Data Analysis

Diversity index (H'), similarity index (E), and dominance (C) are searched based on the formula. To obtain more accurate results used PCA (Begon, 2006; Legendre & Legendre, 1998) using SAS 9.1.3 package.

The diversity index is calculated using the Shannon-Wiener formula (H')

$$H' = -\sum P_i \ln P_i \quad (1)$$

Similarity Index (E)

The Similarity index is known by using the formula: $E = H' / H_{\max}$ (2)

Dominance Index

Dominance is known by using the formula:

$$C = \sum (n_i / N)^2 \quad (3)$$

Results and Discussion

Six genera gastropods found in the Babura river, namely, (1) Pomacea, (2) Filopaludina, (3) Lymnaea, (4) Terebia, (5) Melanoides, (6) Thiara and 1 sub genera, namely Clea (Anentome) (Table. 1). (Narayan *et al*, 2023).

Table 1. Classification of Gastropods in the Babura River to genera/sub-genera

Class	Ordo	Family	Genus	Sub genus	Species	
Gastropoda	Architaenio	Ampulridae	Pomacea		<i>Pomacea</i>	
	Glossa				<i>Canaliculata</i>	
			Viviparidae	Filopaludina		<i>Filopaludina Javanica</i>
	Higropila	Lymnaeidae	Lymnaea		<i>Lymnaea Rubiginosa</i>	
					<i>Terebia granifera</i>	
	Caenogastropoda	Thiaridae	Terebia		<i>Melanoides Tuberculata</i>	
				Melanoides		<i>Thiara scraba</i>
					Thiara	
	Neogastropoda	Nassariidae	Clea		<i>Clea (Anentome) Helena</i>	
				(Anentome)		Neogastropoda

Freshwater gastropods have 18 species from 14 genera and nine families, including four non-native species (Ng *et al*, 2017).

There are no molluscs that dominate on the Kabung island (Kushadiwijayanto *et al*, 2022).

Table 2. Diversity Index (H'), Similarity Indeks (E), and Dominance (D) of Gastropos at Sampling Station

No	Genera/sub genera	Sampling						Total	Average
		I	II	III	IV	V	VI		
1.	Pomacea	-	-	-	-	6,00	14,38	20,38	3,40
2.	Filopaludina	-	-	-	-	5,25	2,88	8,13	1,35
3.	Lymnaea	-	-	-	0,38	-	-	0,38	0,06
4.	Tarebia	-	-	67,13	7,88	-	6,50	55,00	9,17
5.	Melanoides	-	-	57,75	3,50	22,50	2,75	86,50	14,42
6.	Thiara	-	-	121,13	-	9,13	-	130,25	21,71
7.	Clea (Anentome)	-	-	-	0,50	-	-	0,50	0,08
	Total ($\sum D_i$)	-	-	246,00	12,25	42,88	26,50	492,00	50,19
	Average	-	-	35,14	1,75	6,13	3,79		7,17
	Diversity Index (H')	-	-	1,03	0,87	1,18	1,13		

Similarity Index (E)	-	-	0,94	0,63	0,85	0,81
Dominance (C)	-	-	0,14	0,48	0,34	0,37

Diversity at station IV is low and at stations II, V, and VI is moderate. Similarity at stations III, IV, V, and VI is high. The dominance at stations III, V, and VI is low and at station IV is moderate.

The substrate in the Babura river consists of sand, silt, and clay, with the percentage composition of the substrate as presented in Table 3.(Faiz *et al.* 2022).


Table 3. Type of Substrate in Babura River

Station	Substrate %			Substrate Type
	Sand	Silt	Clay	
I	86,80	9,7,80	3,29	Sand Clay
II	75,70	18,50	5,80	Sand Clay
III	78,70	11,30	9,90	Sand Clay
IV	6,20	45,80	44,40	Silt Clay
V	66,70	14,30	19,00	Silt Clay
VI	6,50	65,70	27,70	Silt Clay

The physical and chemical parameters contained in each station are presented in Figures 2.

The temperature in the Babura river is in the range of 20⁰C -28⁰C. There is no difference that fluctuates at each station. The

highest temperature is at station IV and the lowest temperature is at station I (a). The depth in the Babura river are 11.6 cm -86.5 cm, the highest depth is at station VI and the lowest is at station I. At III and IV there is a difference in the third sampling, at station V overall there is very little difference, and at station VI there is a slight decrease. in the seventh sampling (b). Current flow are 0.13 m/s-1.11 m/s, the highest current is at station IV and the lowest is at station III. At stations I, II, and III it is almost evenly distributed, at station IV in the second sampling more high, and at station VI there was a difference in current flow for each sampling (c). The turbidity are 0.70 mg/l-20.50 mg/l, at stations I and II it was evenly distributed for each sampling, at stations III to VI it increased in the 2, 3, 4, and 5 samplings. and 8, the highest at station III and the lowest at station I (d). TDS are 25 m/s-145 m/s, the highest TDS at station VI and the lowest is at station I. At station I there is a slight difference in each sampling, at station II it is very high in sampling 4, at station III there is only a slight difference, at stations IV, V, and VI there is an increase in sampling 1, 3, and 7 (e).



Figures 2. The physical and chemical parameters contained in each station

a. Diagram of water temperature ($^{\circ}\text{C}$), b. Diagram of water depth (cm), c. Diagram of current flow (m/dtk), d. Diagram turbidity (mg/l), e. Diagram, TDS (mg/l), f. Diagram pH, g. Diagram DO (mg/l), h. Diagram BOD (mg/l), i. Diagram Nitrate (mg/l), j. Diagram Phosphate (mg/l).

The pH are 5.5 to 6.93, each sampling is slightly different, the highest pH is at station II and the lowest is at station III (f). DO are 0.40 mg/l-8.12 mg/l, at stations III, IV, and V the difference was large, and at station VI the difference was striking. At stations I and II the DO was > 2 mg/l, at station V at the 8 sampling and at station IV at the third, 4, and 5 sampling the DO < 2 mg/l, the highest DO is at station I and the lowest at station VI (f). The highest BOD is at station IV and the lowest is at station I (g). The nitrates are 0.10 mg/l-3.10 mg/l, the highest at station IV and the lowest at station II. At station I it is slightly different for each sampling, at station II it is almost evenly distributed, at stations III, IV, V, and VI different for each sampling (g). Phosphate was in the range of 0.10 mg/l-2.60 mg/l, the highest phosphate was at station IV and the lowest phosphate was at station II (Tyagi et al. 2023). At stations III to VI there are differences in each sampling (h). The results of testing the hypothesis by species, that in each sampling there is no difference. The results of hypothesis testing based on stations during sampling, the significant difference was only at station III, namely, sampling 1, 3, 4, 5, 6, 7. The results of the hypothesis test of physical and chemical parameters, turbidity and nitrate showed significant differences in taking 1, 2, 3, 4, 5, and 6 Parichelra et al. 2022). The correlation between physical and chemical parameters with the number of species is; (1) temperature, depth, turbidity, DO, nitrate, and phosphate with a very low correlation between the number of species, (2) TDS and BOD with a low correlation between the

number of species, (3) current strength and pH with the number of species have an inverse correlation, which means if the current is high then the number of species is low and if the current is low then the number of species is large (Pramanik et al. 2021).

Relationship Between Each Parameter With Each Species

The test results for each parameter that affect each species, which are very significantly different, namely (1) temperature, depth, TDS to *Pomacea canaliculata*, (2) depth, pH, nitrate to *Filopaludina javanica*, (3) current flow to *Lymnaea rubiginosa*, (4) TDS to *Melanoides tuberculata*, (5) current flow and TDS to *Terebia granifera*, (6) current flow to *Thiara scraba* and *Clea (Anentome)helenae*. While the significant differences were (1) nitrate to *Pomacea canaliculata*, (2) temperature to *Filopaludina javanica*, (3) turbidity to *Lymnaea rubiginosa*, (4) temperature, current flow, turbidity to *Melanoides tuberculata*, (5) depth to *Terebia granifera*, while DO, BOD, and phosphate have no effect. The test results for each parameter that affect each species, which are very significantly different are (1) temperature, depth, TDS to *Pomacea canaliculata*, (2) depth, pH, nitrate to *Filopaludina javanica*, (3) current flow to *Lymnaea rubiginosa*, (4) TDS to *Melanoides tuberculata*, (5) current flow and TDS to *Terebia granifera*, (6) current flow to *Thiara scraba*, and *Clea (Anentome)helenae*, and the significant differences were (1) nitrate to *Pomacea canaliculata*, (2)

temperature to *Filopaludina javanica*, (3) turbidity to *Lymnaea rubiginosa*, (4) temperature, current flow, turbidity to *Melanoides tuberculata*, (5) depth to *Terebia granifera*, while DO, BOD, and phosphate have no effect. *Lymnaea rubiginosa* and *Clea (Anentome)Helena* are found only at station IV so they are markers in the water of Babura river. According to Lomartire *et al*, 2021, the most widely used biomarkers are benthic organisms and fish, which provide a more specific indication of the stressor in the environment Srivastava *et al*. 2022),(Swaney *et al*. 2023).

Principal Component Analysis (PCA)

Parameters affecting gastropod species in the Babura river were then analyzed using PCA

statistics.

The diagram represent




Figure 3. Grouping of Gastropods Based on PCA Analysis

The diagram representing the distribution of gastropods with the influence of physical/chemical parameters on the F1 and F2 axes shows the presence of three groups of gastropods. *Pomacea canaliculata* is affected by TDS, temperature, and depth on the positive F1 axis; *Lymnaea rubiginosa* and *Clea*

ing the distribution of gastropods on the F1 and F2 axes shows the presence of three groups. First, *Pomacea canaliculata* forms a positive F1 axis; second consists of *Filopaludina javanica*, *Lymnaea rubiginosa*, and *Clea (Anentome)helenaf* form the negative F1 axis, and the third *Melanoides tuberculata* and *Terebia granifera* which contributes to the positive F2 axis. Based on the location of the distribution, *Lymnaea rubiginosa* and *Clea (Anentome) helena* have similarities because they form an angle that is $<45^{\circ}$, as well as between *Melanoides tuberculata* and *Terebia granifera*.

(Anentome) helena were affected by currents, flow, *Filopaludina javanica* was affected by nitrate and the pH on the F1 axis was negative, *Melanoides tuberculata* and *Terebia granifera* were affected by turbidity.

PCA results on axis 1 (F1) with an effect of 63.12% and axis 2 (F2) with an effect of 18.59%. The results of PCA on

axis 1 (F1) with a diversity of 42.73% and axis 2 (F2) with a diversity of 23.58 % showed that the population of large and medium-sized lokan clams spread on a

substrate of dust, clay and high organic matter and negatively correlated with conditions pH and temperature (Figure 4).




Figure 4 . Parameters Affecting Gastropods With PCA Analysis

There were 7 species of gastropods found in the Babura river, but not found at stations I and II, because the temperature ranged from 18⁰C-20⁰C. Gastropods were found 8 species in all station, as many as 5 research station with temperature between 25.90⁰C-26.77⁰C (Moghdani *et al*,2013). There are two species, namely *Cassidula nucleus* and *Cassidula angulifera* (Ariyanto *et al* 2018), fewer than those in the Babura River. The larger number is 11 species from 8 families (Normalasari *et al.*, 2019) the highest diversity index is as moderate, the dominance index level is no dominates (Afwanudin *et al*, 2019), 7 gnera, and from 6 families from 4 order (Hecca *et al.*, 2017). According to (Tobing and Harahap (2021). the physical/chemical parameters in the Pandayangan river support gastropod life with 230 individuals consististing of 7 species found. Current flow in the Babura river is slow because the water discharge is small and has many turns but the number of species is small. *Terebia granifera* and *Melanoides tuberculata* were found at

stations III, IV, and VI,. Where as at station V only *Melanoides tuberculata* was found, *Terebia granifera* was not found, in accordance with (Perissinotto *et al.*, 2014) the two species did not coexist. There were three genera found because of their low substrate diversity, which will have an impact on food sources and habitat forb gastropods (Rahmayanti *et al*, 2018). The temperature increase in the present study might be due to the fact that the river segment on that stretch is more open (Chatanga, 2018). DO and water pH are physical and chemical parameters that affect the distribution and diversity of macrozoobenthos with a strong correlation category (0.776) (Dimenta *et al.*, 2020). The statistical analysis with Principal component analysis (PCA) was calculated for measuring the affiliation between environmental variables and number of individuals in sampling sites (Rheca, 2021).

After PCA analysis, continued with multiple regression analysis, presenter in table 4-10.

Table 4. Determinant Coefficient of *Pomacea canaliculata*

No	Regression equation	Determinant coefficient (R ²)
1	$Y = 17,15 - 2,21X_7$	47,40%
2	$Y = 7,78 - 1,35X_7 + 0,10X_2$	53,41%
3	$Y = 9,37 - 1,45X_7 + 0,13X_2 - 0,02X_5$	54,07%
4	$Y = 5,04 - 1,43X_7 + 0,12X_2 - 0,02X_5 + 0,19X_1$	54,39%
5	$Y = 6,11 - 1,43X_7 + 0,12X_2 - 0,02X_5 + 0,13X_1 + 0,02X_4$	54,50%
6	$Y = 6,37 - 1,41X_7 + 0,12X_2 - 0,02X_5 + 0,14X_1 + 0,02X_4 - 1,63X_3$	54,70%
7	$Y = 6,47 - 1,41X_7 + 0,12X_2 - 0,02X_5 + 0,13X_1 + 0,02X_4 - 1,49X_3 - 0,08X_9$	54,72%
8	$Y = -37,70 - 1,41X_7 + 0,10X_2 + 0,00X_5 + 0,50X_1 + 0,01X_4 - 6,29X_3 - 0,03X_9 + 5,63X_6$	61,83%
9	$Y = -46,50 - 0,74X_7 + 0,06X_2 + 0,05X_5 + 0,94X_1 + 0,11X_4 - 2,98X_3 + 0,07X_9 + 4,63X_6 - 0,90X_8$	65,77%
10	$Y = -25,69 - 0,83X_7 + 0,09X_2 + 0,03X_5 + 0,59X_1 + 0,12X_4 + 0,77X_3 + 1,54X_9 + 2,57X_6 - 0,77X_8 - 2,93X_{10}$	69,84%

Table 5. Determinant Coefficient of *Thiara scraba*

No	Regression equation	Determinant coefficient (R ²)
1	$Y = 57,34 - 102,45X_3$	12,82%
2	$Y = 237,36 - 75,27X_3 - 30,19X_6$	17,96%
3	$Y = 65,80 - 119,99X_3 - 5,94X_6 + 5,94X_8$	27,88%
4	$Y = 12,98 - 143,19X_3 + 2,54X_6 + 5,66X_8 + 12,23X_{10}$	31,02%
5	$Y = 58,80 - 146,93X_3 - 13,33X_6 + 5,33X_8 + 15,28X_{10} + 8,82X_7$	40,88%
6	$Y = 59,67 - 147,01X_3 - 13,37X_6 + 5,36X_8 + 15,30X_{10} + 8,77X_7 - 0,00X_5$	40,88%
7	$Y = 58,24 - 147,24X_3 - 13,15X_6 + 5,36X_8 + 15,66X_{10} + 8,77X_7 - 0,00X_5 - 0,34X_9$	40,89%
8	$Y = 49,20 - 158,16X_3 - 8,43X_6 + 7,85X_8 + 16,99X_{10} + 6,54X_7 - 0,12X_5 - 1,00X_9 - 0,70X_4$	42,16%
9	$Y = 144,22 - 155,28X_3 - 12,33X_6 + 8,93X_8 + 14,55X_{10} + 5,12X_7 - 0,13X_5 - 0,19X_9 - 0,65X_4 - 2,71X_1$	42,75%
10	$Y = 152,22 - 154,58X_3 - 12,81X_6 + 9,14X_8 + 14,06X_{10} + 5,19X_7 - 0,16X_5 - 0,07X_9 - 0,67X_4 - 2,99X_1 + 0,08X_2$	42,78%

Table 6. Determinant Coefficient of *Tarebia granifera*

No	Regression equation	Determinant coefficient (R ²)
1	$Y = -2,48 + 2,77X_8$	11,12%
2	$Y = 76,70 + 2,19X_8 - 12,08X_6$	14,19%
3	$Y = 6,86 + 3,49X_8 + 0,94X_6 - 56,40X_3$	24,23%
4	$Y = 23,00 + 3,77X_8 - 1,04X_6 - 56,88X_3 - 0,05X_5$	24,67%
5	$Y = 2,44 + 3,76X_8 + 2,57X_6 - 69,96X_3 - 0,08X_5 + 6,76X_{10}$	28,07%
6	$Y = 2,64 + 3,76X_8 + 2,54X_6 - 69,92X_3 + 0,08X_5 + 6,71X_{10} - 0,04X_9$	28,07%
7	$Y = -22,92 + 2,67X_8 - 0,06X_6 - 67,61X_3 + 0,11X_5 + 7,20X_{10} - 0,32X_9 + 4,81X_7$	33,73%
8	$Y = -26,92 + 3,77X_8 + 2,02X_6 - 72,44X_3 + 0,05X_5 + 7,79X_{10} - 0,61X_9 + 3,82X_7 - 0,31X_4$	34,65%
9	$Y = 32,31 + 4,45X_8 - 0,40X_6 - 70,65X_3 + 0,05X_5 + 6,26X_{10} - 0,11X_9 + 2,93X_7 - 0,27X_4 - 1,69X_1$	35,49%
10	$Y = 17,29 + 4,06X_8 + 0,49X_6 - 71,96X_3 + 0,11X_5 + 7,17X_{10} - 0,34X_9 + 2,79X_7 - 0,23X_4 - 1,18X_1 - 0,15X_2$	35,86%

Table 7. Determinant Coefficient of *Melanoides tuberculata*

No	Regression equation	Determinant coefficient (R ²)
1	$Y = 163,46 - 23,76X_6$	16,46%
2	$Y = 136,76 - 17,38X_6 - 38,51X_3$	23,23%
3	$Y = 42,40 - 4,04X_6 - 63,11X_3 + 3,26X_8$	36,40%
4	$Y = 102,16 - 7,15X_6 - 63,68X_3 + 4,00X_8 - 1,77X_1$	38,07%
5	$Y = 100,42 - 6,97X_6 - 63,94X_3 + 4,05X_8 - 1,75X_1 - 0,02X_4$	38,08%
6	$Y = 109,74 - 10,04X_6 - 65,99X_3 + 4,67X_8 - 0,91X_1 - 0,12X_4 - 0,14X_5$	41,06%
7	$Y = 79,08 - 6,76X_6 - 75,31X_3 + 4,63X_8 - 0,37X_1 - 0,17X_4 - 0,17X_5 + 4,56X_{10}$	42,78%
8	$Y = 86,63 - 7,73X_6 - 74,23X_3 + 4,63X_8 - 0,44X_1 - 0,16X_4 - 0,18X_5 + 3,16X_{10} + 1,27X_9$	42,90%
9	$Y = 58,94 - 6,32X_6 - 76,20X_3 + 3,88X_8 + 0,46X_1 - 0,08X_4 - 0,07X_5 + 4,71X_{10} + 0,87X_9 - 0,25X_2$	44,09%

10	$Y = 51,13 - 6,56X_6 - 75,59X_3 + 3,60X_8 + 0,63X_1 - 0,05X_4 - 0,04X_5 + 4,82X_{10} + 0,82X_9 - 0,25X_2 + 0,64X_7$	44,17%
----	---	--------

Table 8. Determinant Coefficient of *Lymnaea rubiginosa*

No	Regression equation	Determinant coefficient (R2)
1	$Y = -0,28 + 0,99X_3$	34,57%
2	$Y = -0,36 + 0,92X_3 + 0,01X_8$	37,87%
3	$Y = -0,36 + 0,92X_3 + 0,01X_8 - 0,00X_4$	37,94%
4	$Y = -0,36 + 0,96X_3 + 0,02X_8 - 0,00X_4 - 0,04X_{10}$	39,10%
5	$Y = -0,22 + 0,95X_3 + 0,02X_8 - 0,00X_4 - 0,04X_{10} - 0,00X_1$	39,23%
6	$Y = -0,06 + 0,96X_3 + 0,02X_8 - 0,00X_4 - 0,04X_{10} - 0,01X_1 - 0,00X_7$	39,41%
7	$Y = -0,06 + 0,95X_3 + 0,02X_8 - 0,00X_4 - 0,04X_{10} - 0,00X_1 - 0,01X_7 - 0,00X_2$	39,51%
8	$Y = 1,34 + 1,18X_3 + 0,01X_8 + 0,00X_4 - 0,06X_{10} - 0,01X_1 - 0,00X_7 - 0,00X_2 - 0,19X_6$	42,93%
9	$Y = 1,27 + 1,17X_3 + 0,01X_8 + 0,00X_4 - 0,05X_{10} - 0,01X_1 - 0,00X_7 - 0,00X_2 - 0,18X_6 - 0,01X_9$	43,01%
10	$Y = 1,60 + 1,16X_3 + 0,03X_8 - 0,00X_4 - 0,06X_{10} - 0,02X_1 - 0,01X_7 + 0,00X_2 - 0,20X_6 - 0,00X_9 - 0,00X_5$	43,72%

Table 9. Determinant Coefficient of *Filopaludina javanica*

No	Regression equation	Determinant coefficient (R2)
1	$Y = -12,17 + 0,54X_1$	22,10%
2	$Y = -11,61 + 0,51X_1 - 0,005X_2$	22,21%
3	$Y = -11,64 + 0,51X_1 + 0,00X_2 - 0,00X_5$	22,25%
4	$Y = -14,38 + 0,66X_1 - 0,00X_2 + 0,00X_5 - 0,17X_8$	24,10%
5	$Y = -21,80 + 0,79X_1 + 0,00X_2 + 0,01X_5 - 0,31X_8 + 0,50X_7$	28,34%
6	$Y = -22,16 + 0,80X_1 + 0,00X_2 + 0,02X_5 - 0,36X_8 + 0,54X_7 + 0,01X_4$	28,51%
7	$Y = -21,21 + 0,78X_1 + 0,00X_2 + 0,02X_5 - 0,32X_8 + 0,52X_7 + 0,01X_4 - 1,14X_3$	28,96%
8	$Y = -21,88 + 0,81X_1 + 0,00X_2 + 0,01X_5 - 0,34X_8 + 0,52X_7 + 0,01X_4 - 1,53X_3 + 0,29X_9$	30,05%
9	$Y = 36,82 + 0,90X_1 - 0,00X_2 + 0,02X_5 - 0,23X_8 + 0,45X_7 - 0,00X_4 - 3,65X_3 + 0,30X_9 + 2,15X_6$	35,02%
10	$Y = -37,64 + 0,91X_1 - 0,00X_2 + 0,02X_5 - 0,24X_8 + 0,45X_7 - 0,00X_4 - 3,80X_3 + 0,24X_9 + 2,23X_6 + 0,11X_{10}$	35,05%

Table 10. Determinant Coefficient of *Clea (Anentome) helena*

No	Regression equation	Determinant coefficient (R2)
1	$Y = -0,27 + 0,90X_3$	35,47%
2	$Y = -0,31 + 0,87X_3 + 0,008X_8$	36,42%
3	$Y = -0,30 + 0,89X_3 + 0,001X_8 + 0,003X_4$	37,64%
4	$Y = -0,30 + 0,91X_3 + 0,002X_8 + 0,003X_4 - 0,02X_{10}$	37,97%
5	$Y = 0,70 + 1,12X_3 - 0,008X_8 + 0,004X_4 - 0,04X_{10} - 0,16X_6$	41,40%
6	$Y = 0,83 + 1,11X_3 - 0,007X_8 + 0,004X_4 - 0,03X_{10} - 0,17X_6 - 0,001X_2$	41,91%
7	$Y = 0,83 + 1,11X_3 - 0,007X_8 + 0,004X_4 - 0,03X_{10} - 0,17X_6 - 0,00X_2 + 0,001X_7$	41,91%
8	$Y = 0,83 + 1,11X_3 - 0,007X_8 + 0,004X_4 - 0,03X_{10} - 0,17X_6 - 0,00X_2 + 0,00X_7 - 0,00X_5$	41,91%
9	$Y = 1,54 + 1,13X_3 + 0,002X_8 + 0,004X_4 - 0,04X_{10} - 0,20X_6 + 0,00X_2 - 0,007X_7 - 0,00X_5 - 0,02X_1$	43,04%
10	$Y = 1,23 + 1,09X_3 + 0,001X_8 + 0,004X_4 + 0,001X_{10} - 0,17X_6 - 0,00X_2 - 0,006X_7 - 0,00X_5 - 0,01X_1 - 0,04X_9$	44,25%

Description: temperature (X1), depth (X2), current flow (X3), turbidity (X4), TDS (X5), pH (X6), DO (X7), BOD (X8), nitrate (X9), phosphate (X10), number of spesies (Y)

Based on multiple regression analysis, the determinat coefficient of *Pomacea canaliculata* ranged from 14.6%-40.89%, *Thiara scraba* ranged from 12.82%-35.86%, *Tarebia granifera* ranged from 11.12%-35.86%, *Melanoides tuberculata* ranged

from 16.46%-44.17%, *Lymnaea rubiginosa* ranged from 4.57%-43.72%, *Filopaludina javanica* ranged from 22.10%-35.05%, *Clea (Anentome) helena* ranged from 35.47%-44.25%.

From table 4-10, it can be seen that the order of the variable X (physical/chemical parameters) from the highest to the lowest, which can affect each gastropod species found in the Babura river.

Conclusion

Gastropod community in the river consisted of : 6 genera, namely, (1) Pomacea, (2) Filopaludina, (3) Lymnaea, (4) Terebia, (5) Melanoides, (6) Thiara and 1 sub-genera, namely Clea (Anentome). Diversity at station 4 is low and at stations 2, 5, and 6 is moderate. Similarity at stations 3 to 6 is high. The dominance at stations 3, 5, and 6 is low and at station 4 is medium. Parameters that greatly affect the presence of each species of gastropods in the Babura River are: *Pomacea canaliculata* by DO, *Thiara scraba* by current flow, *Tarebia granifera* by BOD, *Melanoides tuberculata* by pH, *Lymnaea rubiginosa* by current flow, *Filopaludina javanica* by temperature, and *Clea (Anentome)helenae* by current flow. The gastropods markers in the Babura river are *Lymnaea rubiginosa* and *Clea (Anentome)helenae*.

Recommendations

We recommend the extend observed station and also the chemical parameters which are possible influence the gastropod properties. We also recommend the other river like the Deli's river in order to know the properties of gastropod.

Ethical clearance

The methodology of sampling caused no harm to the environment and the associated fauna and flora in Babura River.

Funding

No fund has been received for this study.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Albrecht Christian, Stelbrink Björn, Gauffre-Autelin Pauline, Marwoto Ristiyanti M., Rintelen Thomas von, Glaubrecht Matthias. 2020. Diversification of epizoic freshwater limpets in ancient lakes on Sulawesi, Indonesia: Coincidence or co evolution. *Journal of Great Lakes Research*. 46 (2020): 1187-1198
- Abdelhady, A. A., Abdelrahman, E., Elewa, A. M., Fan, J., Zhang, S., & Xiao, J.2018. Phenotypic plasticity of the gastropod *Melanoides tuberculata* in the Nile Delta: A pollution-induced stabilizing selection. *Marine Pollution Bulletin*, 133, 701-710.
<https://doi.org/10.1016/j.marpolbul.2018.06.026>
- Abdel-Satar A.M., Ali M.H., Goher M.E. 2017. Indices of water quality and metal pollution of Nile River, Egypt. *Egyptian Journal of Aquatic Research* 43, 21–29. DOI: 10.1016/j.ejar.2016.12.006
- Afwanudin A., Sarong M A., Efendi R., Deli A., Irham M. 2019. The community structure of Gastropods as bioindicators of water quality in Krueng Aceh, Banda Aceh. *IOP Conf. Series: Earth and Environmental Science* 348(2019) 012122 .
- A. Harahap, P. Hrp, N.K.A.R. Dewi, Macrozoobenthos diversity as an bioindicator of the water quality in the River Kualuh Labuhanbatu Utara, *International Journal of Scientific &*

- Technology Research, 9(4), 2020, pp. 179-183.
doi:10.1088/1755-1315/348/1/012122
- Ariyanto Dafit., Bengen G. Dietriech., Prartono Tri., and WardiatnoYusli. 2018. The association of *Cassidula nucleus* (Gmelin 1791) and *Cassidula angulifera* (Petit 1841) with mangrove in Banggi Coast, Central Java, Indonesia. *AACL Bioflux*. 11 (2). 348 <http://www.bioflux.com.ro/aac>
- Begon M., Townsend C.R., and Harper J.L. 2006. *Ecology: from Individuals to Ecosystems*. Malden, MA: Blackwell Pub.
- Chatanga P., NtuliV., MugomeriE.. 2018. Situational analysis of physico-chemical, biochemical and microbiological quality of water along Mohokare River, Lesotho. *Egyptian Journal of Aquatic Research*, <https://doi.org/10.1016/j.ejar.2018.12.002>
- K.Khairul, R Machrizal, A Harahap,2019. Biological aspects of fish indo pacific tarpon (Megalops cyrinoides Broussonet, 1782) at Belawan River IOP Conference Series: Earth and Environmental Science 348 (1), 012028 vol: | issue : 2019
- Galan Gloria L., Ediza Marilou M., Servasques Marife S., and Porquis Heidi C. 2015. Diversity of Gastropods in the Selected Rivers and Lakes in Bukidnon. *International Journal of Environmental Science and Development*,6(8):615-619.
- Harahap, A, et, all, Macrozoobenthos diversity as anbioindicator of the water quality in the Sungai Kualuh Labuhanbatu Utara, *AACL Bioflux*, 2022, Vol 15, Issue 6.
- Harahap, A.,Budianto Bangun, et all 2022. Analysis Of Water Quality From Bio-Physical-Chemical Factors Of The Asahan River North Sumatra. *Annals of Forest Research*[this link is disabled](#), 2022, 65(1), pp. 1513–1528.
- Hamli H., Azmai S.H. Syed., Hamed Abdul S., Al-Asif Abdulla. 2020. Diversity and Habitat Characteristics of Local Freshwater Gastropoda (Caenogastropoda) from Sarawak, Malaysia. *Singapore Journal of Scientific Research*, 10: 23-27
DOI: 10.3923/sjsres.2020.23.27.
- Hecca Desven., Hidayat Saleh., Dewiyeti Susi. 2017. The Diversity of Water Environment Gastropodain The Water of Empayang-Kasap River in Lahat Regency South Sumatra. *Biovalentia: Biological Research Journal*, 3(1): 1-7
- Jung-Bae Mi & Seuk-Park Young. 2020. Key Determinants of Freshwater Gastropod Diversity and Distribution: The Implications for Conservation and Management. *MDPI*, 12(7):10.3390/w12071908
- Kushadiwijayanto Arie Antasari., Helena Shifa., Warsidah., Apriansyah. 2022. Diversity Of Mollucs (Bivalves And Gastropods) In Kabung Island, West Kalimantan, Indonesia. *Spermonde*, 7(2):34-37.
- Legendre P. & Legendre L. 1998. *Numerical Ecology*. 2nd English edn Elsevier Science BV, Amsterdam.
- Lomartire Silvia., Marques Joao C., Goncalves Ana M.M. 2021. Biomarkers based tools Assess environmental and chemical stressors in aquatic systems. *Ecological*

- Indicators* 122(2021)107201 1-9,
www.elsevier.com/locate/ecolind.
- Moghdani S., Amiri F., Ghanbari F., Saki Entezami M., Tabatabaei T., and Pourkhan M., 2013. Water quality assessment with biological indicators: Mond protected area, Iran. *Journal of Biodiversity and Environmental Sciences*, 3(9):80-89.
- Ng Ting Hui ., Dulipat Jasrul., Foon Junn Kitt., Lopes-Lima Manuel., Zieritz Alexandra, Liew Thr-Seng. 2017. A preliminary checklist of the freshwater snailsof Sabah (Malaysian Borneo) deposited in theBORNEENSIS collection, Universiti Malaysia Sabah. *ZooKeys* 673: 105–123.
- Normalasari., Melani Winny Retna., Apriadi Tri. 2019. Structure Community of Gastropods in A ir Kelubi water, Resun Pesisir Village, Lingga Utara Subdision. *Jurnal Akuatiklestari*, 2(2): 10-19.
- Perissinotto R., Miranda N., Raw J., and Peer N.2014. Biodiversity census of Lake St Lucia, Simangaliso Wetland Park (South Africa): gastropod molluscs. *Zoo Keys*, 440:1–43
- Rahmayanti., Nazira F.K., Dewi K.A., Oktaviani F.D., Millaty K.N.I., Prasetya A.T., Sasmita. B.H., Nashrurrokhman M., Roshitafandi A.D., Febiansi D., Sartika W.H., Zulfikar G.W., Kurnia N.R., Islami A.D., and Tranggono A.Y. 2018. Biodiversity of gastropod in the Sombu Beach, Wakatobi, Indonesia. *The 2nd International Symposium on Marine and Fisheries Research IOP Publishing IOP Conf. Series: Earth and Environmental Science*, 139 (2018) 012013.
- Raw L. Jacqueline., Miranda F. A., Nelson., and PerissinottoRenzo. 2015. Chemical cues released by heterospecific competitors: behaviouralresponses of native and alien invasive aquatic gastropods. *Aquat Sci*, 77:655–666.
- Rekha Kalimuthu., Anbalagan Sankarappan., Dinakaran Sundaram. 2021. Distributional ecology of snails (Gastropoda: Mollusca) in seasonal ponds of Tamil Nadu, South India. *Acta Ecologica Sinica*, 41(2021):410-415
- Shinta Herlina Eka., Utami Purnama Julia., Adiwijaya Saputra. 2020. Potential Stunting in Riverside Peoples (Study on Pahandut Urban Village, Palangka Raya City). *Budapest International Research and Critics InstituteJournal (BIRCI-Journal)*, 3(3): 1618-1625
- Sinambela Masdiana., Barus Ternala Alexander., Manurung Binari.,Wahyuningsih Hesti. 2019. Gastropods Community in Babura River, Medan city. *IOP Conf. Series: Earth and Environmental Science* 305 (2019) 012092 IOP Publishing
- Harahap A, Mahadewi EP, Ahmadi D, Tj HW, Ganiem LM, Rafika M, Hartanto A. 2021. Monitoring of macroinvertebrates along streams of Bilah River, North Sumatra, Indonesia. *Intl J Conserv Sci* 12 (1): 247- 258.
- Narayan, Vipul, et al. "Enhance-Net: An Approach to Boost the Performance of Deep Learning Model Based on Real-Time Medical Images." *Journal of Sensors* 2023 (2023).
- NARAYAN, VIPUL, A. K. Daniel, and Pooja Chaturvedi. "FGWOA: An Efficient Heuristic for Cluster Head

- Selection in WSN using Fuzzy based Grey Wolf Optimization Algorithm." (2022).
- Babu, S. Z., et al. "Abridgement of Business Data Drilling with the Natural Selection and Recasting Breakthrough: Drill Data With GA." Authors Profile Tarun Danti Dey is doing Bachelor in LAW from Chittagong Independent University, Bangladesh. Her research discipline is business intelligence, LAW, and Computational thinking. She has done 3 (2020).
- Narayan, Vipul, A. K. Daniel, and Pooja Chaturvedi. "E-FEERP: Enhanced Fuzzy based Energy Efficient Routing Protocol for Wireless Sensor Network." *Wireless Personal Communications* (2023): 1-28.
- Faiz, Mohammad, et al. "Improved Homomorphic Encryption for Security in Cloud using Particle Swarm Optimization." *Journal of Pharmaceutical Negative Results* (2022): 4761-4771.
- Tyagi, Lalit Kumar, et al. "Energy Efficient Routing Protocol Using Next Cluster Head Selection Process In Two-Level Hierarchy For Wireless Sensor Network." *Journal of Pharmaceutical Negative Results* (2023): 665-676.
- Paricherla, Mutyalaiah, et al. "Towards Development of Machine Learning Framework for Enhancing Security in Internet of Things." *Security and Communication Networks* 2022 (2022).
- Pramanik, Sabyasachi, et al. "A novel approach using steganography and cryptography in business intelligence." *Integration Challenges for Analytics, Business Intelligence, and Data Mining*. IGI Global, 2021. 192-217.
- Srivastava, Swapnita, et al. "An Ensemble Learning Approach For Chronic Kidney Disease Classification." *Journal of Pharmaceutical Negative Results* (2022): 2401-2409.
- Sawhney, Rahul, et al. "A comparative assessment of artificial intelligence models used for early prediction and evaluation of chronic kidney disease." *Decision Analytics Journal* 6 (2023): 100169.