

ISSN 2063-5346



DETECTION OF VIRULENCE FACTORS OF PSEUDOMONAS AERUGINOSA ISOLATED FROM CLINICAL SAMPLES

Raghad Z. Suleiman^{1*}, Bashar sadeq noomi^{2*}, Ibrahim Omar Saeed^{3*}

Article History: Received: 10.05.2023

Revised: 29.05.2023

Accepted: 09.06.2023

Abstract

The aim of this research is to study the effectiveness of *Pseudomonas aeruginosa* bacteria isolated from clinical samples to produce Protease, Lipase, Amylase, Urease, DNase and Hemolysine. One hundred samples were collected from patients hospitalized during July 2020 till the end of January 2021 to isolate and identify *P. aeruginosa* from clinical sources including infections of wounds and burns. Twenty three isolates were found to be belonging to *P. aeruginosa*, which mean an isolate rate of (23%) of the total samples. The diagnosis was carried out based on phenotypic characteristics and biochemical tests. *P. aeruginosa* was found in (10) out of (38) wound infections samples, (13) out of (62) burn infections samples. This means that isolates from the wound infections constituted the highest percentage of (26%) of the total samples, while the lowest percentage of (20.9%) was obtained from the burn infections samples. The results showed that (24.7%) of total isolates were Hemolysin producer on blood agar, It was also found that (60.8%) of total isolates were Amylase producer on Starch agar, (69.5%) of total isolates were Protease and Lipase producer on Skim milk agar and on Rhan agar, (21.7%) of total isolates were Urease producer on Urea agar and (30.4%) were DNase producer on a DNase agar medium.

Keywords: *Pseudomonas aeruginosa*, Clinical Samples, virulence factors.

¹Dept. of Biology - College of Science - Tikrit University - Tikrit - Iraq

²College of Veterinary Medicine Verified, University of Tikrit, Salah Al-Din, Iraq.

³Department of Biology, College of Science, University of Tikrit, Salah Al-Din, Iraq.

*Corresponding Author, Raghadzeyad99@yahoo.com

* Dr.ibrahim1977@tu.edu.iq

DOI:10.48047/ecb/2023.12.9.140

Introduction

P. aeruginosa is an important opportunistic pathogen causing life-threatening acute infections in individuals with compromised immune systems. It is also the most common cause of chronic mental infections and skin infections, and the main cause of morbidity and death in patients with hereditary cirrhosis (Kostylev *et al.*, 2019).

Pseudomonas Gram-negative bacteria appear as short chains or single cells, or may be straight or curved bacilli, moving by a single polar flagellum. *P. aeruginosa* are endemic in different environments such as soil, water, on surfaces, etc. (Chevallereau, 2017).

Pseudomonas transmitted mainly by dust-borne, tap water and bed sheets. Moreover, patients with chronic skin lesion are considered as a source for others lying in the same hall, as well as the operating theater bed, surgical instruments, floors, walls, cupboards, lighting, anesthesia device, ventilation and sterilizers. Doctors, nurses, other staff and visitors also take part in the process of transmitting these bacteria by direct contact or by using some multi-use uniforms (Nikbin *et al.*, 2012; Inglisa *et al.*, 2009).

Diseases caused by *P. aeruginosa* bacteria depend on their possession of multiple virulence factors that enable them to break down tissues and invade the bloodstream. Among the most important of these factors are Protease, DNase and Hemolysine (Wilhel *et al.*, 1999; Ernst, 1999).

Hospital infection is one of the main problems facing health care staff, especially those who deal with serious surgical cases as a result of bacterial contamination in the surgical operating theaters. This in turn leads to the spread of hospital infections among patients too (Iiyama *et al.*, 2017). The spread of these infections has many dangerous effects, including an increase in the morbidity and

mortality rate, the prolonged stay of the patient in the hospital and the increased need to use antibiotics that have an effective effect on the bacteria (Finnan *et al.*, 2004).

Materials and Methods

1. Samples Collection

The samples of this study were collected from clinical sources using cotton swab under supervision of a specialist physician. Then each sample was activated on a nutrient agar, incubated, taken with a tube and be carefully taken out of the tube so as not to touch the tube wall, and finally transplanted on a special plate for the growth of *P. aeruginosa* for the purpose of purification.

It is isolated from patients hospitalized, (100) clinical samples from two places of the patients hospitalized. The sample collection period lasted from the end of July (2020) to the end of January (2021).

2. Preparation of Culture Media

Readymade culture media were prepared according to the manufacturer's instructions. Three types of media were prepared as follows:

2.1 Blood Agar

This medium was prepared according to (Rutala W.A., 1996).

2.2 Cetrimide Agar

This medium was prepared according to (Holt *et al.*, 1984).

2.3 Skim Milk Agar

This medium was prepared according to (Kalai, 2009).

3. Isolation of *P. aeruginosa*

After the samples were collected by swabs, they were cultured by loops using the spreader method on blood agar and MacConkey agar. The plates were incubated at 37°C for 24 hours. After the incubation period, the results are read in which *P. aeruginosa* was found not to

ferment lactose sugar. Then, the developing bacterial colonies were transferred to *Pseudomonas* agar medium and the dishes were incubated at 37°C for 24 hours. The isolates carrying the characteristics of *P. aeruginosa* were selected and re-cultivated on a Cetrimide agar, which is regarded as a special medium for *P. aeruginosa*, and then incubated at 37°C for 24 hours. These colonies were also purified more than once to obtain good, pure isolates for the purpose of diagnosis confirmatory of *P. aeruginosa*.

4. Diagnosis of Bacterial Isolates

Bacterial isolates were diagnosed based on cultural and microscopic characteristics as well as biochemical tests as follows:

4.1 Cultural Characteristics

In order to study the agronomic characteristics of the bacterial isolates, the bacterial isolates ability to grow was tested at first on a group of culture media, including *Pseudomonas* agar, special *Pseudomonas* agar, Cetrimide agar, and selective *P. aeruginosa* according to (Brooks *et. al.*, 2007). They were also cultured on MacConkey agar and Blood base agar in order to diagnose cultural characteristics in terms of colony color and shape.

4.2 Microscopic Examination

A microscopic examination of the developing bacterial cells was carried out and stained with a gram stain solutions according to (Brooks *et. al.*, 2007).

4.3 Biochemical Tests

In order to diagnose the isolates at the species level, the purified colonies were subjected to two types of biochemical tests, Oxidase test and Catalase test According to (Rahman, 2006).

5. Determination of Virulence Factors

5.1 Hemolysin Production

According to (Wisplinghoff, 2017) , the ability of the isolates to produce bacterial hemolysin and ability of *P. aeruginosa* bacteria to analyze blood.

5.2 Protease Production

According to (Cruickshank *et al.*, 1975), the ability of the isolates to produce bacterial Protease and ability of *P. aeruginosa* bacteria to analyze protein .

5.3 DNase Production

According to (Heidi *et. al.*, 2010), the ability of the isolates to produce bacterial DNase and ability of *P. aeruginosa* bacteria to hydrolyze DNA .

5.4 Lipase Production

According to (Ghafil & Hassan, 2014), the ability of the isolates to produce bacterial Lipase and ability of *P. aeruginosa* bacteria degradation of lipids .

5.5 Amylase Production

According to (Sivaramakrishnan, 2007), the ability of the isolates to produce bacterial Amylase and ability of *P. aeruginosa* bacteria to hydrolysis starch .

5.6 Urease Production

According to (Qassem, 2006), the ability of the isolates to produce bacterial Urease and ability of *P. aeruginosa* bacteria to hydrolysis urea (ammonia).

Results and Discussion

1. Isolation and Identification

The results of isolation and diagnosis of patients hospitalized samples showed the presence of bacterial growth belonging to the species *P. aeruginosa* in (23) out of (100) samples of clinical sources including (10) out of (38) isolates from the Wound infections, and(13) out of (62) isolates from the Burns infections , as listed in Table (1).

Table 1: Clinical *P. aeruginosa* isolates according to sources

| Type of samples | Total clinical samples | <u><i>P. aeruginosa</i></u> isolates | Percentages |
|------------------|------------------------|--------------------------------------|-------------|
| Wound infections | 38 | 10 | 26% |
| Burns infections | 62 | 13 | 20.9% |
| Total | 100 | 23 | 23% |

2. Agricultural Characteristics


2.1 Growth on Cetrimide Agar

It appears in the form of green colonies as shown in Fig. (1).

**Figure 1:** *P. aeruginosa* green colonies grown on Cetrimide medium

2.2 Growth on Nutrient Agar

It produce (80%) pyocyanin when grown on Nutrient agar medium. Large colonies with little convexity and flat edges were observed as shown in Fig. (2).

**Figure 2:** *P. aeruginosa* colonies grown on Nutrient medium

2.3 Growth on Blood Agar

Transparent halo appeared around the colonies that were grown on a stimulating enrichment medium as shown in Fig. (3).



Figure 3: *P. aeruginosa* colonies grown on Blood medium

2.4 Growth on Skim Milk agar

The ability of the isolates to produce protease enzyme was investigated using Skim Milk agar medium as shown in Fig. (4).




Figure 4: *P. aeruginosa* colonies grown on Skim Milk agar

2.5 Production of the deoxyribonucleic acid (DNase) enzyme

The results of this research showed the ability of the isolates to produce this enzyme as shown in Fig. (5). Transparent halo appeared around the bacterial colonies cultivated on a DNase agar medium after adding HCL.



Figure 5: Production of the deoxyribonucleic acid (DNase) enzyme

2.6 Production of the Lipase enzyme

The ability of the isolates to produce Lipase enzyme was investigated using Rhan agar medium as shown in Fig. (6).




Figure 6: Production of the Lipase enzyme on Rhan agar medium

2.7 Production of the Amylase enzyme

The ability of the isolates to produce Amylase enzyme was investigated using Starch agar medium as shown in Fig. (7).




Figure 7: Production of the Amylase enzyme on Starch agar

2.8 Production of the Urease enzyme

The ability of the isolates to produce Urease enzyme was investigated using Urea agar base as shown in Fig. (8).




Figure 8: Production of the Urease enzyme on Urea agar

Virulence factors

The results showed that (24.7%) of total isolates were Hemolysin producer on blood agar, It was also found that (69.5%) of total isolates were Protease producer on Skim milk agar, (30.4%) were DNase producer, (69.5%) were Lipase producer, (60.8%) were Amylase producer, (21.7%) were Urease producer, Table (2)

Table (2) shows a summary of the Virulence factors found in this research in the *P. aeruginosa* isolates out of 23 Clinical samples.

Table 2: *P. aeruginosa* isolates out of 23 Clinical samples

| Virulence factors | No. | percentage |
|-------------------|-----|------------|
| Hemolysin | 8 | 24.7%% |
| Protease | 16 | 69.5%% |
| DNase | 7 | 30.4%% |
| Lipase | 16 | 69.5% |
| Amylase | 14 | 60.8% |
| Urease | 5 | 21.7% |

References

- Kostylev M., Kim D.Y., Smalley N.E., Salukhe I., Greenberg E.P. and Dandekar A.A. (2019). Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy. Proceedings of the National Academy of Sciences, 116(14), 7027-7032.
- Chevallereau A. (2017). Comprehensive study of new virulent bacteriophages: from Transcriptomic and Mechanistic Characterisations towards Evolutionary Perspectives. Thesis, Université Sorbonne, Paris. English.
- Nikbin V., Aslani N. and Ebrahimipour K. (2012). Molecular identification and detection of virulence genes among pseudomonas aeruginosa isolated from different infectious origins. Iran Microbial 4(3): 118-123.
- Inglisa F., Gardnab A. and Cornelis P. (2009). Spite and virulence in the bacterium *P. aeruginosa*. PNAS Journal 106(14): 5703-5707.
- Wilhelm S., Tommassen J. and Jaeger K.E. (1999). A novel lipolytic enzyme located in the outer membrane of *Pseudomonas aeruginosa*. Journal of Bacteriology, 181(22): 6977-6986.
- Ernst R.K., Yi E.C., Guo L., Lim K.B., Burns J.L., Hackett M. and Miller S.I. (1999). Specific lipopolysaccharide found in cystic fibrosis airway *Pseudomonas aeruginosa*. Science, 286: 1561-1565.
- Liyama K., Takahashi E., Lee J.M., Mon H., Morishita M., Kusakabe T. and Yasunaga-Aoki C. (2017). Alkaline protease contributes to pyocyanin production in *Pseudomonas aeruginosa*. FEMS Microbiology Letters, 364(7), fnx051.
- Finnan S., Morrissey J.P., O'Gara F. and Boyd E.F. (2004). Genome diversity of *Pseudomonas aeruginosa* isolates from cystic fibrosis patients and the hospital environment", Journal of Clinical Microbiology, 42(12): 5783-5792.
- Rutala W.A. (1996). APIC guideline for selection and use of disinfectants. Amer J. Infect. Control 24(4): 313-342.
- Holt J.G., Krieg N.R., Sneath P.H., Staley J.T. and Williams S.T. (1994). Bergy's manual of determinative bacteriology. 9th Edition, Williams and Wilkins, 1063.
- Kalai M., Oana, C., Claus S., Sternberg P. and Peter, W. (2009). Mucoid conversion of *Pseudomonas aeruginosa* by hydrogen peroxide: a mechanism for virulence activation

- in the cystic fibrosis lung. Clin. Infect. Dis., 23: 1109-16.
12. Brooks G.F., Butel J.S. and Morse S.A., (2007). Jawetz, Melnick and Adelberg's Medical Microbiology, 24th Edition, McGraw-Hill.
 13. Rahman F.M. (2006). Genetic analysis of protease from *Pseudomonas aeruginosa* isolated from different human infections. MSc Thesis. College of Science, University of Babylon.
 14. Wisplinghoff H. (2017). *Pseudomonas* spp., *Acinetobacter* spp. and miscellaneous gram-negative bacilli in infectious diseases. Elsevier, pp. 1579-1599.
 15. Cruickshank R., Duguid J.P. and Swain R.H.A. (1975). Medical Microbiology. 12th Edition, London and New York .Churchill Livingstone.
 16. Heidi M., Laetitia Ch.M. and Shawn L. (2010). Environmental Microbiology. J. of Biology, Medicine: 10.1111/j.1462-2920x.
 17. Al-Tikrity A.L. (2009). Bacteriological and genetical study of *Pseudomonas aeruginosa* isolated from different human infection. MSc Thesis, College of Science, Tikrit University.
 18. Al-Shuwaikh R.M.A. (2006). Production and characterization of diseases from isolated *Pseudomonas aeruginosa* bacteria and its relationship to some antibiotics. PhD Thesis, College of Science, Al-Mustansiriya University.
 19. Al-Doori L.M.I. and Maarouf M.N. (2018). Isolation and identification of types of common bacteria from different infections and investigation of some factors of their virulence. Al-Rafidain Science Journal, Microbiology 27(5): pp. 99-111.
 20. Al-Doori M.N. (2009). Genetic and molecular study of some vancomycin-resistant gram-positive *Cocci* isolated from Tikrit city. PhD Thesis, College of Education, Tikrit University.
 21. Al-Tamimi A.A.H. (2000). Isolation and characterization of the sticky layer of adherent *Staphylococcus epidermidis*. MSc Thesis, College of Science, Baghdad University.
 22. Sultan N.M.M. (2001). Diagnostic and genetic study of *Coagulase-negative Staphylococcus* isolated from cases of urinary tract infection in puberty women. MSc Thesis, College of Science, Mosul University.
 23. Ghafil J.A. and Hassan SH.S. (2014). Effect of cultural conditions on lipase production *Pseudomonas aeruginosa* isolated from Iraqi soil. World J. Experm. Biosci., 2(1):13-18.
 24. Sivaramakrishnan, S., Gangadharan, D., Nampoothiri, K. M., Soccol, C. R., & Pandey, A. "Alpha amylase production by *Aspergillus oryzae* employing solid-state fermentation". Journal of scientific and industrial research, 66 (8), 621, (2007). Sivaramakrishnan, (2007).
 25. Qassem, K. W. (2006). The Effect of Some Chemical and Physical Factors on the Sensitivity of Locally Isolated *Pseudomonas aeruginosa* to Antibiotics. M. Sc. Thesis, College of Science, University of Baghdad.