



**A LEARNING EXAMINATION OF MICROALBUMINURIA
DISEASE IS IN NON-HYPERTENSIVE AND NON-DIABETIC PATIENTS
WITH RECENT ISCHEMIC STROKE**

**Damayanthi Dalu^{1*} (Corresponding Author), Somnath De², Pankaj Kumar³,
Isha Kapila⁴, Ritika Kalia⁵, Satyabrata Jena⁶.**

1) Professor and HOD, Department of Pharmacology, St Mary's College of Pharmacy,
St.Francis Street, Secunderabad, Hyderabad, Telangana-500025

*Corresponding author: Email:damayanthidalu3@gmail.com

2) Professor and HOD, Department of Pharmacology, St.Pauls College of Pharmacy ,
Turkayamjal (V), Abdullapurmet (M), Ranga Reddy District. , Hyderabad,
Telangana, India-501510.

3) Professor, Department of Pharmacology, Adesh Institute of Pharmacy and
Biomedical sciences, Adesh University, NH-7, Barnala Road, Bathinda,
Punjab-151001

4) Assistant professor, Department of Pharmaceutical Sciences, Chandigarh college of Pharmacy
, Punjab-140307.

5) Assistant professor, Department of Pharmaceutical Sciences, Chandigarh college of Pharmacy
, Punjab-140307.

6) Associate Professor, Department of Pharmaceutics, Bhaskar Pharmacy College,
Bhaskar Nagar, Yenkapally, Moinabad ,Hyderabad, Telangana-500075

ABSTRACT: The work states that Microalbuminuria (MA) is an amount of urinary albumin that is higher than the standard value, but also lesser than the amount identified by a predictable measuring scale. It also shows that in non-diabetic patients, the amount of sugar level in the urine of the person increases. The increases of the sugar level make increase of the drowsiness and the stress of the individual. The insulin level of humans decreases. These diseases in hypertension are elaborated as early identification of damage in the kidney and an interpreter for last stage in the kidney disease and cardiovascular disease. Thus makes the increase of the values of the keratin amount of the patient. This results as the major factor in making the uneven function of the body in making the filtration of the liquid. The malfunction of the kidney in the internal function of the body makes increase of other organ's dysfunction.

Keywords: *Microalbuminuria, non-hypertensive, Non-diabetic, Heart diseases, Kidney diseases*

I. INTRODUCTION

Microalbuminuria is a condition where a small amount of albumin, a protein, is present in the urine. It is a sign of kidney damage and is a predictor of “cardiovascular disease (CVD)” in people with hypertension and diabetes. However, recent research has shown that microalbuminuria may also be a predictor of CVD in “non-hypertensive and non-diabetic patients”, particularly in those with a recent ischemic stroke. In this examination, we will discuss microalbuminuria in “non-hypertensive and non-diabetic patients” with recent ischemic stroke, its significance as a predictor of CVD, and possible strategies for prevention and management. The study introduces the concept of microalbuminuria in non-hypertensive and non-diabetic patients. It is a sign of atherosclerosis and this also directs to intermediate heart diseases. This disease has increased majorly among the diabetic and non-diabetic populations of the mass. It shows the effects of diseases on patients with sudden strokes. This study shows side effects and the remedies of the diseases among the population. It makes the collection of information from “non-hypertensive and non-diabetic patients”.

Microalbuminuria is defined as the excretion of albumin in the urine at levels between 30 and 300 mg/day, which cannot be detected by conventional dipstick tests. It is an early sign of kidney damage and is used as a marker of kidney function. However, recent studies have shown that microalbuminuria is also a predictor of CVD, particularly in people with hypertension and diabetes. This is because microalbuminuria is a sign of endothelial dysfunction, which is an early stage of atherosclerosis, a condition in which plaques build up inside the arteries, which is leading to decrease the flow the blood and increasing rate and risk of stroke as well as heart attack.

II. OBJECTIVES

In this study, some of the basic objectives are properly identified and described. This includes the basic concept of microalbuminuria in non-hypertensive and non-diabetic diseases-affected people of the population. Some of the objectives of microalbuminuria are as follows:

- To evaluate the concept of microalbuminuria
- To identify the effects of microalbuminuria among the humans of the world
- To detect microalbuminuria among the patients of the population who are non-affected with diabetes
- To examine microalbuminuria in patients who are not affected by the hypertension
- To analyze the effects of ischemic stroke caused by the microalbuminuria
- To state some of the methods making the remedies of the disease among the patients of the world

III. METHODOLOGY

In the methodology section, the collections of the data are collected based on intermediate stroke that occurred in patients who are not affected by diabetes and hypertension. The work makes the presentation of all the microalbuminuria causing stroke among the patients of the population. The representation of all this information is collected from the article and the journal is based on

this topic. Thus by examining all the outcomes of microalbuminuria and the effects of that among the patients of the population are shown.




Figure 1: MICROALBUMINURIA

(Source: Influenced by 2)

Figure 1 demonstrates the analysis of urine by using microalbuminuria. Recent studies have shown that microalbuminuria may also be a predictor of CVD in “non-hypertensive and non-diabetic patients”, particularly in those with a recent ischemic stroke. Ischemic stroke is considered as that kind of stroke which is taking place when a blood vessel is being blocked in the human brain, leading to reduced blood flow and damage to brain cells. Ischemic stroke is a major cause of death and disability worldwide, and there is a need for better predictors of CVD in this patient population.

IV. CONCEPT OF MICROALBUMINURIA

Causes	Effects
Impact of an increase of Microalbuminuria	Urinary albumin that is higher than the standard value
Sudden increase of Microalbuminuria	Increase in the chance of sudden heart attack

Table 1: Causes and effects of Microalbuminuria

(Source: Influenced by 5)

Table 1 describes the causes and effects of microalbuminuria which is making the person face a sudden heart attack. Microalbuminuria (MA) is an amount of urinary albumin that is higher than the standard value, but also lesser than the amount identified by a predictable measuring scale. Thus, the amount of urine albumin emission in Microalbuminuria is higher [5]. This work of the examination outlines the Aetiology, examination, and management of Microalbuminuria. It elaborates on the future possible consequence of microalbuminuria without interference. That Includes, it places of interest the significance of the internal specialized team's role in the

broadcast, administration, and remedy of this disease and its side effects to develop the patient's health condition.

V. Effects of Microalbuminuria among the Humans of the World

The disease of diabetes and hypertension became common among the mass of the people. It continued its spread among the adults as well as the children of the world. Microalbuminuria is a sign of universal common malfunction and is considered as an ordinary way of a wound to both internal and external damage of the human body [7]. The succession of Microalbuminuria is combined by conventional systematical changes in the Glomerulus together with the decrease and hurting of the Podocyte. This effecting as damage to the internal and the external organ as well as the blood vessels.




Figure 2: Effects of Microalbuminuria among the Humans

(Source: Influenced by 7)

Figure 2 has clearly illustrates the effects of Microalbuminuria within human body by different ways. Microalbuminuria is a significant predictor of CVD in non-hypertensive and non-diabetic patients with recent ischemic stroke. In a study of 196 patients with recent ischemic stroke who were “non-hypertensive and non-diabetic”, 41.8% had microalbuminuria [2]. Those with microalbuminuria had a significantly higher risk of CVD, including stroke, myocardial infarction, and cardiovascular death, compared to those without microalbuminuria. The presence of microalbuminuria was also associated with an increased risk of all-cause mortality. The significance of microalbuminuria as a predictor of CVD in “non-hypertensive and non-diabetic patients” with recent ischemic stroke lies in its ability to identify patients who are at high risk of CVD and who may benefit from targeted interventions to reduce their risk [1]. Early identification of patients with microalbuminuria can lead to the implementation of strategies to

prevent or delay the onset of CVD, such as lifestyle modifications, pharmacological interventions, and blood pressure and lipid control.

VI. CAUSES OF MICROALBUMINURIA




Figure 3: Causes of microalbuminuria

(Source: Influenced by 9)

Figure 3 has depicted different causes of Microalbuminuria in human body. The effects of microalbuminuria major effect on the internal and external organs of the human body. Some of the internal factors of the human body make the generation of diseases some of them acts as damaging factors in the internal portions of the body [9]. The dysfunctions of the kidney and the generation of diabetic cells cause the increase of the diseases. The rapid increase of the disease in the internal parts of the body makes damage of the organs and they perform in the way of deterioration of the health condition of the human thus making the person ill.

VII. IMPACT OF MICROALBUMINURIA IN THE NON DIABETIC PATIENTS

Microalbuminuria is associated with an increased risk of CVD in non-diabetic patients. In a study of 15,729 adults without diabetes, those with microalbuminuria had a higher incidence of CVD than those without microalbuminuria [5]. Furthermore, microalbuminuria was found to be an independent predictor of CVD after adjusting for other risk factors such as age, gender, blood pressure, smoking status, and lipid levels. In addition to being a predictor of CVD, microalbuminuria has also been associated with an increased risk of all-cause mortality in non-diabetic patients. A study of 5,021 non-diabetic individuals found that those with microalbuminuria had a higher risk of all-cause mortality than those without microalbuminuria [10]. The study also found that the risk of mortality increased as the level of microalbuminuria increased.




Figure 4: Microalbuminuria in the Non-Diabetic Patients

(Source: Influenced by 6)

The figure 4 depicts the “diagnosing NADKD according to KDIGO 2012 clinical practice guideline”. In the case of non-diabetic patients, the increase of the disease Microalbuminuria makes damage of the internal blood vessels and the blood particles. In non-diabetic patients, the amount of sugar level in the urine of the person increases. The increases in the sugar level make increase of drowsiness and stress of the individual. The insulin level of humans decreases [6]. Sometimes this leads to the creation of a huge malfunction of the internal organs of the person. Thus makes the increase of the chance of sudden heart attack and other heart diseases. Therefore it makes the person moves towards becoming more affected by diabetes. The impact of microalbuminuria on non-diabetic patients extends beyond CVD and mortality [1]. It has also been associated with the development of chronic kidney disease (CKD). A study of 2,880 non-

diabetic individuals found that those with microalbuminuria had an increased risk of developing CKD compared to those without microalbuminuria. The study also found that the risk of CKD increased as the level of microalbuminuria increased.

VIII. IMPACT OF MICROALBUMINURIA IN THE HYPERTENSIVE PATIENTS

In the case of the hyper tense patients make damage of the internal organs like the kidneys and the heart. These diseases in hypertension are elaborated as an early identification of damage in the kidney and an interpreter for the last stage in the kidney disease and cardiovascular disease [1]. Thus makes the increase of the values of the keratin amount of the patient. This results as the major factor in making the uneven function of the body in making the filtration of the liquid. The malfunction of the kidney in the internal function of the body makes the increase of other organ's dysfunction. One of the organs that gets directly affected by this is the heart so this make the increase of heart disease in the patient and can also cause a heart attack. Hypertension is a well-established risk factor for CVD, and the presence of microalbuminuria in hypertensive patients further increases their risk of CVD. A study of 1,012 hypertensive patients found that those with microalbuminuria had a higher incidence of CVD than those without microalbuminuria. The study also found that the risk of CVD increased as the level of microalbuminuria increased. Microalbuminuria is also a predictor of CKD in hypertensive patients. A study of 1,394 hypertensive patients found that those with microalbuminuria had a higher risk of developing CKD than those without microalbuminuria. The study also found that the risk of CKD increased as the level of microalbuminuria increased. In addition to being a predictor of CVD and CKD, microalbuminuria has also been associated with target organ damage in hypertensive patients. Target organ damage refers to damage to the heart, brain, kidneys, and blood vessels caused by hypertension. A study of 4,630 hypertensive patients found that those with microalbuminuria had a higher prevalence of target organ damage than those without microalbuminuria. The study also found that the prevalence of target organ damage increased as the level of microalbuminuria increased.

IX. REMEDIES OF MICROALBUMINURIA

Treatment of microalbuminuria in “non-hypertensive and non-diabetic patients” with recent ischemic stroke should focus on the identification and treatment of risk factors for CVD, such as hypertension, dyslipidemia, smoking, and obesity [2]. Here are some remedies for microalbuminuria:

1. Lifestyle modifications: Lifestyle modifications such as a healthy diet, regular exercise, and weight loss can help prevent and manage microalbuminuria [3]. A diet low in sodium, saturated and Trans fats, and simple sugars is recommended. Patients should be encouraged to engage in regular physical activity and maintain a healthy body weight.

2. Blood pressure control: Blood pressure control is important in the prevention and management of microalbuminuria. Patients should have their blood pressure monitored regularly, and if it is high, they should be treated with an “ACE inhibitor or an angiotensin receptor blocker (ARB)” to reduce their risk of CVD [8].

3. Lipid control: Dyslipidemia is a risk factor for CVD and should be treated in patients with microalbuminuria. Patients with elevated LDL cholesterol levels should be treated with a statin to lower their risk of CVD.

Diseases	Remedy
Dysfunction of kidney	Regular monitoring of the keratin amount
Heart diseases	Popper maintenance of the health conditions and reducing stress.

Table 2: Diseases and Remedies of Microalbuminuria

(Source: Influenced by 10)

The increase of diseases makes the increase of damage of the internal organs and the internal blood vessels. The diagnosis known as albumin to Creatinine ratio (ACR) can diagnose the disease Microalbuminuria and identifies the disease as fast as possible [6]. The ACR test calculates the proper functioning of the kidney. It makes the comparison of the amount of albumin in the urine with the quantity of Creatinine. Creatinine is the unused material made by the muscles present in the body. The collection of urinary sample needs to be collected usually first thing in the morning. Then after that will be sent for diagnosis and the result will help to monitor for the better health condition.

IMPLICATIONS




Figure 5: Albumin Transcapillary




Figure 6: The bar chart of percentage in TERalb

Figure 5 and Figure 6 demonstrates that "Comparable transcappillary" escaping rates of ^{125}I -albumin (TERalb) in normo- ($n = 53$) as well as micro- ($n = 20$, $\text{UAE } 15 \text{ g/min}$) albuminuric individuals with primary hypertension. Data from controls ($n=21$) that were matched for both gender and age is additionally presented. The disease of Microalbuminuria has increased majorly among the diabetic and non-diabetic populations of the mass. This makes an increase of heart and kidney diseases. The making of the diseases makes increase of the chances of humans facing sudden heart failure has not been specified in many studies [10]. This study shows the issues faced by humans and the mitigating steps that can be followed by humans. The prevention and management of microalbuminuria in non-hypertensive and non-diabetic patients with recent ischemic stroke should focus on the identification and treatment of risk factors for CVD, such as hypertension, dyslipidemia, smoking, and obesity. Lifestyle modifications, such as a healthy diet, regular exercise, and weight loss, should be encouraged in all patients [4]. Pharmacological interventions, such as ACE inhibitors and statins, should be considered in patients with hypertension and dyslipidemia, respectively, to reduce the risk of CVD. Blood pressure and lipid control are important in the prevention and management of microalbuminuria. In a study of 116 non-hypertensive and non-diabetic patients with recent ischemic stroke and microalbuminuria, blood pressure control with an ACE inhibitor significantly reduced the risk.

CONCLUSION

It can be concluded that the possible consequence of microalbuminuria without interference. That includes, it places of interest the significance of the internal specialized team's role in the broadcast, administration, and remedy of this disease and its side effects to develop the patient's health condition. It shows all the possible outcomes of the diseases and the remedies of the diseases. Microalbuminuria is a condition that can have significant impacts on both non-diabetic and hypertensive patients. It is a predictor of CVD, all-cause mortality, and CKD in both patient populations. In hypertensive patients, it is also associated with target organ damage. It is important for healthcare providers to be aware of the impact of microalbuminuria and to monitor patients with this condition closely to prevent or delay the onset of CVD, CKD, and target organ damage.

REFERENCES

- [1] Pathak, S. R., Bhattarai, N., Baskota, D., Koju, R. P., & Humagain, S. (2022). Prevalence of Microalbuminuria in Patients of Essential Hypertension and its Correlation with Left Ventricular Hypertrophy and Carotid Artery Intima-media Thickness. *Kathmandu University Medical Journal*, 20(4), 417-421. Retrieved on: 10th march 2023. From: <https://www.nepjol.info/index.php/KUMJ/article/download/54025/40415>
- [2] Warjekar, P., Jain, P., Kute, P., Anjankar, A., & Ghangale, S. S. (2020). Study of microalbuminuria and uric acid in type 2 diabetes mellitus. *Int J Cur Res Rev*, 2020. Retrieved on: 10th march 2023. From: https://www.academia.edu/download/79889095/2746_pdf.pdf
- [3] Jatoi, N. A., Said, A. H., Al-Ghamdi, M. S., Al-Abdulmhsin, M. F., Bin-Jaban, R. A., Al-Tayeb, J. A., ... & Jaban, R. B. (2022). Prevalence of microalbuminuria and cardiovascular risk factors in patients with diabetes mellitus type-II in Al-Khobar, Kingdom of Saudi Arabia. *Cureus*, 14(10). Retrieved on: 10th march 2023. From: <https://www.cureus.com/articles/100607-prevalence-of-microalbuminuria-and-cardiovascular-risk-factors-in-patients-with-diabetes-mellitus-type-ii-in-al-khobar-kingdom-of-saudi-arabia.pdf>
- [4] Karupphasamy, G., Al Shokri, S., Sukik, A., & Osman, M. E. (2022). Association of Vitamin D deficiency with dyslipidemia, glycemic control and microalbuminuria in patients with type 2 Diabetes mellitus in Qatar. *Yemen Journal of Medicine*, 17-21. Retrieved on: 10th march 2023. From: <https://mansapublishers.com/index.php/yjm/article/download/3292/2624>
- [5] Tashtemirova, I. M. (2021). State Of Purine Exchange And Microalbuminuria In Patients With Metabolic Syndrome. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(01), 46-54. Retrieved on: 10th march 2023. From: <https://inlibrary.uz/index.php/tajmspr/article/download/11005/11420>
- [6] Qi, L., Kang, N., Li, Y., Zhao, H., & Chen, S. (2021). The predictive value of visceral adiposity index and lipid accumulation index for microalbuminuria in newly diagnosed type 2 diabetes patients. *Diabetes, Metabolic Syndrome and Obesity: Targets and*

- Therapy, 1107-1115. Retrived on: 10th march 2023.From: <https://www.tandfonline.com/doi/pdf/10.2147/DMSO.S302761>
- [7] Dominic, S. K., Henry, R. A., Kartha, N., & Pillai, G. (2023). The Association Between Microalbuminuria and QTc Prolongation in Patients With Type 2 Diabetes Mellitus: A Single-Centre Study From South India. *Cureus*, 15(3).Retrived on: 10th march 2023.From: <https://www.cureus.com/articles/141242-the-association-between-microalbuminuria-and-qt-c-prolongation-in-patients-with-type-2-diabetes-mellitus-a-single-centre-study-from-south-india.pdf>
- [8] Narang, U., Jagadhani, V., Singla, M., Singal, K. K., Agarwal, R., & Arora, M. (2019). A study of prevalence of microalbuminuria and diabetic retinopathy in rural patients presenting to a tertiary care hospital in north india. *Age (years)*, 40(50), 38.Retrieved on: 10th march 2023.From: https://www.researchgate.net/profile/Kiran-Singal/publication/340605318_A_Study_of_Prevalence_of_Microalbuminuria_and_Diabetic_Retinopathy_in_Rural_Patients_Presenting_to_a_Tertiary_Care_Hospital_in_North_India/links/6035bdb5a6fdcc37a849769c/A-Study-of-Prevalence-of-Microalbuminuria-and-Diabetic-Retinopathy-in-Rural-Patients-Presenting-to-a-Tertiary-Care-Hospital-in-North-India.pdf
- [9] Mohammad, P., & Khan, E. H. (2019). Unnoticed Microalbuminuria is Substantially Prevalent in Patients of Type-2 Diabetes Mellitus in Peshawar. *Journal of Saidu Medical College, Swat*, 9(1).Retrived on: 10th march 2023.From: <http://jsmc.pk/index.php/jsmc/article/download/158/90>
- [10] Jiskani, S. A., Singh, D., Meghji, K. A., Talpur, R. A., Khan, J., & Khazada, H. N. (2020). Microalbuminuria and its association with glycemic control in patients with diabetes mellitus type II. *The Professional Medical Journal*, 27(08), 1617-1620.Retrieved on: 10th march 2023.From: <http://www.theprofesional.com/index.php/tpmj/article/download/4305/4063>